

Intersection bodies and valuations

Monika Ludwig*

Dedicated to Prof. Erwin Lutwak on the occasion of his sixtieth birthday

Abstract

All $GL(n)$ covariant star-body-valued valuations on convex polytopes are completely classified. It is shown that there is a unique non-trivial such valuation. This valuation turns out to be the so called ‘intersection operator’– an operator that played a critical role in the solution of the Busemann-Petty problem.

2000 AMS subject classification: 52A20 (52B11, 52B45)

A function Z defined on the set \mathcal{K} of convex bodies (that is, of convex compact sets) in \mathbb{R}^n or on a certain subset \mathcal{C} of \mathcal{K} and taking values in an abelian semigroup is called a *valuation* if

$$ZK + ZL = Z(K \cup L) + Z(K \cap L),$$

whenever $K, L, K \cup L, K \cap L \in \mathcal{C}$. *Real valued* valuations are classical and Blaschke obtained the first classification of such valuations that are $SL(n)$ invariant in the 1930s. This was greatly extended by Hadwiger in his famous classification of continuous, rigid motion invariant valuations and characterization of elementary mixed volumes. See [13], [17], [32], [33] for information on the classical theory and [1]–[4], [15], [16], [25], [26], [28] for some of the more recent results.

In [24], [27], a classification of *convex-body-valued* valuations $Z : \mathcal{P} \rightarrow \mathcal{K}$ was obtained where \mathcal{P} is the set of convex polytopes in \mathbb{R}^n containing the origin and addition in \mathcal{K} is Minkowski addition of convex bodies (defined by $K + L = \{x + y : x \in K, y \in L\}$). A valuation Z is called $GL(n)$ *covariant*, if there exists a $q \in \mathbb{R}$ such that for all $\phi \in GL(n)$ and all bodies K ,

$$Z(\phi K) = |\det \phi|^q \phi ZK,$$

where $\det \phi$ is the determinant of ϕ . It is called $GL(n)$ *contravariant*, if there exists a $q \in \mathbb{R}$ such that $\phi \in GL(n)$ and all bodies K

$$Z(\phi K) = |\det \phi|^q \phi^{-t} ZK,$$

where ϕ^{-t} is the transpose of the inverse of ϕ . Since each body $K \in \mathcal{K}$ is determined by its support function, $h(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$, where $h(K, u) = \max\{u \cdot x : x \in K\}$ and where $u \cdot x$ denotes the standard inner product of u and x , these valuations can be defined via support functions. For $n > 2$, the classification theorems [27] are the following. An

*Research was supported in part by the European Network PHD, MCRN-511953

operator $Z : \mathcal{P} \rightarrow \mathcal{K}$ is a $\text{GL}(n)$ contravariant valuation if and only if there is a constant $c \geq 0$ such that

$$Z P = c \Pi P$$

for every $P \in \mathcal{P}$. Here ΠP is the projection body of P , that is, $h(\Pi P, u) = \text{vol}(P|u^\perp)$ for $u \in S^{n-1}$, where vol is the $(n-1)$ -dimensional volume, u^\perp is the subspace orthogonal to u , and $P|u^\perp$ is the image of the orthogonal projection of P onto u^\perp . An operator $Z : \mathcal{P} \rightarrow \mathcal{K}$ is a non-trivial $\text{GL}(n)$ covariant valuation if and only if there are constants $c_0 \in \mathbb{R}$ and $c_1 \geq 0$ such that

$$Z P = c_0 m(P) + c_1 M P$$

for every $P \in \mathcal{P}$. Here an operator is called *trivial*, if it is a linear combination of the identity and central reflection, while $m(P)$ is the moment vector and $M P$ is the moment body of P , defined by,

$$m(P) = \int_P x \, dx \quad \text{and} \quad h(M P, u) = \int_P |x \cdot u| \, dx,$$

for $u \in S^{n-1}$.

These results establish a classification of $\text{GL}(n)$ covariant and contravariant valuations within the Brunn-Minkowski theory. In this paper we ask the corresponding question in the dual Brunn-Minkowski theory. In the dual theory convex bodies are replaced by star bodies and Minkowski addition of convex bodies is replaced by radial addition of star bodies (see next section for definitions). The natural question to ask is for a classification of *star-body-valued* valuations.

Let \mathcal{S} denote the set of star bodies in \mathbb{R}^n , where a set $K \subset \mathbb{R}^n$ is a star body, if it is sharshaped with respect to the origin and has a continuous radial function $\rho(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$ (defined by $\rho(K, u) u \in \partial K$). Let \mathcal{P}_0 denote the set of convex polytopes in \mathbb{R}^n that contain the origin in their interiors and let $P^* = \{x \in \mathbb{R}^n : x \cdot y \leq 1 \text{ for every } y \in P\}$ denote the polar body of $P \in \mathcal{P}_0$.

Theorem. *An operator $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ is a non-trivial $\text{GL}(n)$ covariant valuation if and only if there is a constant $c \geq 0$ such that*

$$Z P = c I P^*$$

for every $P \in \mathcal{P}_0$.

Here $I P^*$ is the *intersection body* of $P^* \in \mathcal{P}_0$, that is, the star body whose radial function is given for $u \in S^{n-1}$ by

$$\rho(I P^*, u) = \text{vol}(P^* \cap u^\perp).$$

In recent years, these intersection bodies have attracted increased interest within different subjects. They first appear in Busemann's [5] theory of area in Finsler spaces and were first explicitly defined and named by Lutwak [29]. Intersection bodies turned out to be critical for the solution of the *Busemann-Petty problem*: If the central hyperplane sections of an origin-symmetric convex body in \mathbb{R}^n are always smaller in volume than those of another such body, is its volume also smaller? Lutwak [29] showed that the answer to the Busemann-Petty problem is affirmative if the body with the smaller sections

is an intersection body of a star body. This led to the final solution that the answer is affirmative if $n \leq 4$ and negative otherwise (see [7], [8], [10], [18], [19], [20], [35], [38], [39]). For further applications of intersection bodies, see [6], [11], [12], [14], [21], [34], and the books and surveys [9], [22], [23], [31], [36], [37].

The next section lists some basics regarding convex bodies, star bodies and valuations. Section 2 contains the proof of the theorem.

1 Notation and background material

General references on convex bodies and star bodies are the books by Gardner [9], Leichtweiß [23], Schneider [36], and Thompson [37]. We work in Euclidean n -space, \mathbb{R}^n , and write $x = (x_1, \dots, x_n)$ for $x \in \mathbb{R}^n$. Let e_1, \dots, e_n denote the vectors of the standard basis of \mathbb{R}^n . For $x, y \in \mathbb{R}^n$, let $x \cdot y = x_1 y_1 + \dots + x_n y_n$ denote the inner product of x and y and let $|x|$ denote the length of x .

Let $K \in \mathcal{S}$. Then its radial function can be extended to $v \in \mathbb{R}^n$, $v \neq 0$, by

$$\rho(K, v) = \max\{\lambda \geq 0 : \lambda v \in K\}.$$

It follows immediately that for $s > 0$ and $\phi \in \text{GL}(n)$,

$$\rho(K, sv) = \frac{1}{s} \rho(K, v) \quad \text{and} \quad \rho(\phi K, v) = \rho(K, \phi^{-1}v). \quad (1)$$

The radial sum $K_1 \tilde{+} K_2$ of $K_1, K_2 \in \mathcal{S}$ is the star body whose radial function is given by

$$\rho(K_1 \tilde{+} K_2, v) = \rho(K_1, v) + \rho(K_2, v).$$

The set \mathcal{S} equipped with the operation $\tilde{+}$ is an abelian semigroup and $\{0\}$ is its neutral element.

Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ *contravariant of weight* $q \in \mathbb{R}$, that is, for all $\phi \in \text{GL}(n)$ and all $P \in \mathcal{P}_0$,

$$Z \phi P = |\det \phi|^q \phi^{-t} Z P.$$

We associate with Z an operator $Z^* : \mathcal{P}_0 \rightarrow \mathcal{S}$ by setting $Z^* P = Z P^*$ for $P \in \mathcal{P}_0$. Let $P_1, P_2, P_1 \cup P_2 \in \mathcal{P}_0$. Since

$$(P_1 \cup P_2)^* = P_1^* \cap P_2^* \quad \text{and} \quad (P_1 \cap P_2)^* = P_1^* \cup P_2^*,$$

we obtain

$$\begin{aligned} Z^* P_1 \tilde{+} Z^* P_2 &= Z P_1^* \tilde{+} Z P_2^* &= Z(P_1^* \cup P_2^*) \tilde{+} Z(P_1^* \cap P_2^*) \\ &= Z(P_1 \cap P_2)^* \tilde{+} Z(P_1 \cup P_2)^* &= Z^*(P_1 \cap P_2) \tilde{+} Z^*(P_1 \cup P_2). \end{aligned}$$

Thus Z^* is a valuation on \mathcal{P}_0 . Let $P \in \mathcal{P}_0$ and $\phi \in \text{GL}(n)$. Since

$$(\phi P)^* = \phi^{-t} P^* \quad (2)$$

and since Z is $\text{GL}(n)$ *contravariant of weight* q , we obtain

$$Z^*(\phi P) = Z(\phi P)^* = Z(\phi^{-t} P^*) = |\det \phi|^{-q} \phi Z^* P. \quad (3)$$

Thus Z^* is $\text{GL}(n)$ *covariant of weight* $-q$.

2 Proof of the Theorem

Lutwak [30] showed that for all $\phi \in \text{GL}(n)$ and all $K \in \mathcal{S}$

$$I(\phi K) = |\det \phi| \phi^{-t} I K.$$

By (3), $P \mapsto I P^*$ is a $\text{GL}(n)$ covariant valuation on \mathcal{P}_0 and we prove that up to multiplication with a constant this is the unique non-trivial such valuation. The proof consists of three steps. First, we extend $\text{GL}(n)$ covariant valuations defined on \mathcal{P}_0 to valuations defined on a larger set of polytopes. Next, we derive a classification of valuations which are $\text{GL}(n)$ covariant of weight $q \geq 0$. Then we derive a classification of valuations which are $\text{GL}(n)$ contravariant of weight $q > 0$. This classification of contravariant valuations and (3) provide a classification of valuations which are $\text{GL}(n)$ covariant of weight $q < 0$. Combined these results prove the theorem.

2.1 Extension

Let $\overline{\mathcal{P}}_0$ denote the set of convex polytopes P which are either in \mathcal{P}_0 or are the intersection of a polytope $P_0 \in \mathcal{P}_0$ and a polyhedral cone with apex at the origin and at most n facets. As a first step, we extend valuations $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ to simple valuations on $\overline{\mathcal{P}}_0$. Here a valuation is called *simple* if $Z(P) = \{0\}$ for every $P \in \overline{\mathcal{P}}_0$ with dimension less than n .

We need the following definitions. For $A, A_1, \dots, A_k \subset \mathbb{R}^n$, let $[A_1, \dots, A_k]$ denote the convex hull of A_1, \dots, A_k and let

$$A^\perp = \{x \in \mathbb{R}^n : x \cdot y = 0 \text{ for every } y \in A\}.$$

For a central hyperplane H (that is, a hyperplane containing the origin), let H^+ and H^- denote the complementary closed halfspaces bounded by H . Let $\mathcal{P}_0(H)$ denote the set of convex polytopes in H that contain the origin in their interiors relative to H .

Let $\mathcal{C}_+(S^{n-1})$ denote the set of non-negative continuous functions on the unit sphere S^{n-1} and let $\overline{\mathcal{C}}_+(S^{n-1})$ denote the set of non-negative functions that are continuous almost everywhere on S^{n-1} . Note that if $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ is a valuation then the operator Y defined by $Y P(\cdot) = \rho(Z P, \cdot)$ is a valuation taking values in $\mathcal{C}_+(S^{n-1})$. Let H be a central hyperplane and let $A \subset S^{n-1}$. For $P \in \mathcal{P}_0(H)$, we say that $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ is *vanishing on A at P* if

$$\lim_{u, v \rightarrow 0} Y[P, u, v] = 0 \text{ locally uniformly on } A$$

for $u \in H^- \setminus H, v \in H^+ \setminus H$. For $P \in \mathcal{P}_0(H)$, we say that Y is *bounded at P* if there exists a constant $c \in \mathbb{R}$ such that

$$Y[P, u, v](x) \leq c$$

for every $x \in S^{n-1}$ and $u \in H^- \setminus H, v \in H^+ \setminus H$ if $|u|, |v| \leq 1$ and $[P, u, v] = [P, u] \cup [P, v]$.

Lemma 1. *Let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be a valuation such that for every central hyperplane H and $P \in \mathcal{P}_0(H)$, Y is bounded and vanishing on $S^{n-1} \setminus H$ at P . Then Y can be extended to a simple valuation $Y : \overline{\mathcal{P}}_0 \rightarrow \overline{\mathcal{C}}_+(S^{n-1})$ and for $P \in \overline{\mathcal{P}}_0$ bounded by central hyperplanes H_1, \dots, H_n , $Y P$ is continuous and bounded on $S^{n-1} \setminus (H_1 \cup \dots \cup H_n)$.*

Proof. Let \mathcal{P}_j , $j = 1, \dots, n$, be the set of convex polytopes P such that there exist $P_0 \in \mathcal{P}_0$ and hyperplanes H_1, \dots, H_j containing the origin with linearly independent normal vectors and

$$P = P_0 \cap H_1^+ \cap \dots \cap H_j^+. \quad (4)$$

Define Y on \mathcal{P}_j , $j = 1, \dots, n$, inductively, starting with $j = 1$, in the following way. For $P \in \mathcal{P}_j$ and $u \in H_1 \cap \dots \cap H_{j-1}$, $u \in H_j^- \setminus H_j$, set

$$Y P = \lim_{u \rightarrow 0} Y[P, u] \quad \text{on} \quad S^{n-1} \setminus (H_1 \cup \dots \cup H_j). \quad (5)$$

We show that Y is well defined (that is, the limit in (5) exists and does not depend on the choice of H_j), that $Y P$ is continuous and bounded on $S^{n-1} \setminus (H_1 \cup \dots \cup H_j)$, that for every hyperplane H and $P \in \mathcal{P}_j(H)$, we have for $u, v \in H_1 \cap \dots \cap H_j$, $u \in H^+ \setminus H$, $v \in H^- \setminus H$,

$$\lim_{u, v \rightarrow 0} Y[P, u, v] = 0 \quad \text{locally uniformly on} \quad S^{n-1} \setminus H \quad (6)$$

and

$$Y[P, u, v] \text{ is uniformly bounded on } S^{n-1} \text{ for } |u|, |v| \leq 1, \quad (7)$$

and that Y has the following additivity properties:

If $P \in \mathcal{P}_{j-1}$ and H is a hyperplane such that $P \cap H^+, P \cap H^- \in \mathcal{P}_j$, then

$$Y P = Y(P \cap H^+) + Y(P \cap H^-) \quad \text{on} \quad S^{n-1} \setminus (H_1 \cup \dots \cup H_{j-1} \cup H). \quad (8)$$

If $P, Q, P \cap Q, P \cup Q \in \mathcal{P}_j$ are defined by (4) with halfspaces H_1^+, \dots, H_j^+ , then

$$Y P + Y Q = Y(P \cup Q) + Y(P \cap Q) \quad \text{on} \quad S^{n-1} \setminus (H_1 \cup \dots \cup H_j). \quad (9)$$

The operator Y is well defined and a valuation on \mathcal{P}_0 . Suppose that Y is well defined by (5) on \mathcal{P}_{k-1} , that $Y P$ is continuous and bounded on $S^{n-1} \setminus (H_1 \cup \dots \cup H_{k-1})$ for $P \in \mathcal{P}_{k-1}$ and that (6), (7), (8) (if $k > 1$) and (9) hold for $j < k$.

First, we show that the limit in (5) exists and that for $P \in \mathcal{P}_k$ bounded by hyperplanes H_1, \dots, H_k , $Y P$ is bounded and continuous on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$. Let $u' \in H_1 \cap \dots \cap H_{k-1}$, $u' \in H_k^- \setminus H_k$ be chosen such that $[P, u] \subseteq [P, u']$ and $-u' \in P$. Then applying (9) with $j = k - 1$ gives on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$Y[P, u] + Y[P \cap H_k, u', -u'] = Y[P, u'] + Y[P \cap H_k, u, -u']. \quad (10)$$

Consequently, for $x \in S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$|Y[P, u](x) - Y[P, u'](x)| \leq Y[P \cap H_k, u', -u'](x) + Y[P \cap H_k, u, -u'](x). \quad (11)$$

Combined with (6) for $j = k - 1$, this implies that the limit in (5) exists locally uniformly and that $Y P$ is continuous on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$. By (10) we have

$$Y[P, u'](x) \leq Y[P, u](x) + Y[P \cap H_k, u', -u'](x).$$

For u fixed, $Y[P, u]$ is bounded by the induction assumption. Let $u' \rightarrow 0$. Then (7) implies that $Y P$ is bounded on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$. For $k > 1$ we show that $Y P$ as defined by (5) does not depend on the choice of the hyperplane H_k . Let $u \in H_1 \cap \dots \cap H_{k-1}$,

$u \in H_k^- \setminus H_k$. Choose $w \in H_2 \cap \dots \cap H_k$, $w \in H_1^- \setminus H_1$. Then applying (8) for $j = k - 2$ gives on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$Y[P, u, w] = Y([P, u, w] \cap H_k^+) + Y([P, u, w] \cap H_k^-). \quad (12)$$

We have $[P, u, w] \cap H_k^- = [P \cap H_k, u, w]$ and $w \in H_k$. By (5) and (6) for $j = k - 2$, we get on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$\lim_{u \rightarrow 0} Y[P \cap H_k, u, w] = 0.$$

Combined with $[P, u, w] \cap H_k^+ = [P, w]$, this implies that on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$\lim_{u \rightarrow 0} Y[P, u, w] = Y[P, w]. \quad (13)$$

Similarly, we get on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$\lim_{w \rightarrow 0} Y[P, u, w] = Y[P, u]. \quad (14)$$

Note that by an argument similar to (11) $\lim_{u, w \rightarrow 0} Y[P, u, w]$ exists. Thus (12) combined with (13) and (14) implies that

$$\lim_{u, w \rightarrow 0} Y[P, u, w] = \lim_{u \rightarrow 0} Y[P, u] = \lim_{w \rightarrow 0} Y[P, w] = YP. \quad (15)$$

Thus Y is well defined on \mathcal{P}_k .

Next, we show that (6) and (7) hold for $j = k < n$. Let $\varepsilon > 0$ be chosen. Let $P \in \mathcal{P}_k(H)$ be bounded by H_1, \dots, H_k . Since $P \subset H$, YP is defined on $S^{n-1} \setminus H$. Let $x \in S^{n-1} \setminus H$. Choose $z \in H \cap H_1 \cap \dots \cap H_{k-1}$ and $z \in H_k^- \setminus H_k$. Then $[P, z] \in \mathcal{P}_{k-1}(H)$ and by (6) for $j = k - 1$,

$$Y[P, z, u, v] < \varepsilon \quad (16)$$

locally around x for $u, v \in H_1 \cap \dots \cap H_k$, $u \in H^- \setminus H$ and $v \in H^+ \setminus H$ with $|u|, |v|$ sufficiently small. Since $[P \cap H_k, u, v] \in \mathcal{P}_{k-1}(H_k)$, (6) for $j = k - 1$ implies that

$$\lim_{w \rightarrow 0} Y[P \cap H_k, u, v, -w, w] = 0 \quad \text{locally uniformly} \quad (17)$$

for $w \in H \cap H_1 \cap \dots \cap H_{k-1}$ and $w \in H_k^- \setminus H_k$. Since Y is a valuation,

$$Y[P, z, u, v] + Y[P \cap H_k, -w, w, u, v] = Y[P, w, u, v] + Y[P \cap H_k, -w, z, u, v].$$

Let $w \rightarrow 0$, then by (5) and (17)

$$Y[P, z, u, v] = Y[P, u, v] + Y[P \cap H_k, z, u, v]. \quad (18)$$

Since $Y \geq 0$, (18) combined with (16) implies that

$$Y[P, u, v] \leq \varepsilon$$

locally around x for $|u|, |v|$ sufficiently small. Thus (6) holds for $j = k$. It follows from (7) that $Y[P, z, u, v]$ is uniformly bounded for $|u|, |v| \leq 1$. Thus (18) implies that (7) holds for $j = k$.

Next, we show that (8) holds for $j = k$. Let $P \in \mathcal{P}_{k-1}$, that is, there exist $P_0 \in \mathcal{P}_0$ and hyperplanes H_1, \dots, H_{k-1} such that $P = P_0 \cap H_1^+ \cap \dots \cap H_{k-1}^+$. Choose $u \in H_1 \cap \dots \cap H_{k-1}$, such that $u \in P \cap H^+ \setminus H$ and $-u \in P \cap H^-$. Then $P, [P \cap H, u, -u], [P \cap H^+, -u], [P \cap H^-, u]$ have the hyperplanes H_1, \dots, H_{k-1} in common. Applying (9) for $j = k - 1$ gives on $S^{n-1} \setminus (H_1 \cup \dots \cup H_{k-1} \cup H)$

$$Y P + Y[P \cap H, u, -u] = Y[P \cap H^+, -u] + Y[P \cap H^-, u].$$

By (6) and definition (5), this implies that (8) holds for $j = k$.

Finally, we show that (9) holds for $j = k$. Choose $u \in H_1 \cap \dots \cap H_{k-1}$, $u \notin H_k$ such that $-u \in P \cap Q$. Applying (9) for $j = k - 1$ shows that on $S^{n-1} \setminus (H_1 \cup \dots \cup H_k)$

$$Y[P, u] + Y[Q, u] = Y[P \cup Q, u] + Y[P \cap Q, u].$$

Because of definition (5) this implies that (9) holds for $j = k$.

The induction is now complete and Y is extended to $\overline{\mathcal{P}}_0$. As last step, we show that Y is a valuation on $\overline{\mathcal{P}}_0$. In addition to (8) and (9) it suffices to prove that if $P \in \mathcal{P}_n$ and H is a hyperplane such that $P \cap H^+, P \cap H^- \in \mathcal{P}_n$, then

$$Y P = Y(P \cap H^+) + Y(P \cap H^-) \quad (19)$$

on $S^{n-1} \setminus (H_1 \cup \dots \cup H_n \cup H)$.

First, let $n = 2$. Let P be bounded by H_1, H_2 , and let $P \cap H^+$ and $P \cap H^-$ be bounded by H_1, H and H, H_2 , respectively. For $u \in H \cap (H_1^- \setminus H_1) \cap (H_2^- \setminus H_2)$, it follows from (8) that $Y[P, u] = Y[P \cap H^+, u] + Y[P \cap H^-, u]$. By (5), this implies that

$$\lim_{u \rightarrow 0} Y[P, u] = Y(P \cap H^+) + Y(P \cap H^-). \quad (20)$$

On the other hand, it follows from (8) that

$$\begin{aligned} Y[P, u] &= Y([P, u] \cap H_1^+) + Y([P, u] \cap H_1^-) \\ &= Y[P, w] + Y([P, u] \cap H_1^- \cap H^-) + Y([P, u] \cap H_1^- \cap H^+) \\ &= Y[P, w] + Y[P \cap H_1, u] + Y[0, u, w]. \end{aligned} \quad (21)$$

where $w \in H_1$ depends on u . Because of (5), we have $\lim_{u \rightarrow 0} Y[P, w] = Y P$ and because of (6), we have $\lim_{u \rightarrow 0} Y[P \cap H_1, u] = 0$. By (8), $Y[P \cap H_2, u] = Y[P \cap H_2, w] + Y[0, u, w]$. Since by (6) $\lim_{u \rightarrow 0} Y[P \cap H_2, u] = 0$ and $\lim_{u \rightarrow 0} Y[P \cap H_2, w] = 0$, this implies that $\lim_{u \rightarrow 0} Y[0, u, w] = 0$. Thus it follows from (21) that $\lim_{u \rightarrow 0} Y[P, u] = Y P$. Combined with (20) this implies (19).

Second, let $n \geq 3$. Let $P = P_0 \cap H_1^+ \cap \dots \cap H_n^+$, $P_0 \in \mathcal{P}_0$. Since $P \cap H^+, P \cap H^- \in \mathcal{P}_n$, we can say that $P \cap H^+$ is bounded by H_1, H, H_3, \dots, H_n and that $P \cap H^-$ is bounded by H, H_2, H_3, \dots, H_n , where $H_1 \cap H_2 \cap \dots \cap H_{n-1} \subseteq H$. Therefore on $S^{n-1} \setminus (H_1 \cup \dots \cup H_n \cup H)$

$$Y P = \lim_{u \rightarrow 0} Y[P, u]$$

and

$$Y(P \cap H^+) = \lim_{u \rightarrow 0} Y[P \cap H^+, u], \quad Y(P \cap H^-) = \lim_{u \rightarrow 0} Y[P \cap H^-, u]$$

where $u \in H_1 \cap H_2 \cap \dots \cap H_{n-1}$, $u \in H_n^- \setminus H_n$. Applying (8) for $j = n$ shows that

$$Y[P, u] = Y[P \cap H^+, u] + Y[P \cap H^-, u].$$

Because of definition (5) this implies (19). This completes the proof of the lemma. \square

We also require the following lemmas. The proofs are similar to that of Lemma 1 and are omitted.

Lemma 2. *Let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be a valuation such that for every central hyperplane H , Y is vanishing on S^{n-1} at $\mathcal{P}_0(H)$. Then Y can be extended to a simple valuation $Y : \overline{\mathcal{P}}_0 \rightarrow \mathcal{C}_+(S^{n-1})$.*

Lemma 3. *Let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be a valuation such that for every central hyperplane H , Y is vanishing and bounded on $S^{n-1} \setminus H^\perp$ at $\mathcal{P}_0(H)$. Then Y can be extended to a simple valuation $Y : \overline{\mathcal{P}}_0 \rightarrow \overline{\mathcal{C}}_+(S^{n-1})$ and for $P \in \overline{\mathcal{P}}_0$ bounded by hyperplanes H_1, \dots, H_n , $Y P$ is continuous and bounded on $S^{n-1} \setminus (H_1^\perp \cup \dots \cup H_n^\perp)$.*

Lemma 4. *Let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be a valuation such that for every central hyperplane H , Y is vanishing on $S^{n-1} \setminus H^\perp$ at $\mathcal{P}_0(H)$. Then Y can be extended to a simple valuation $Y : \overline{\mathcal{P}}_0 \rightarrow \overline{\mathcal{C}}_+(S^{n-1})$ and for $P \in \overline{\mathcal{P}}_0$ bounded by hyperplanes H_1, \dots, H_n , $Y P$ is continuous on $S^{n-1} \setminus (H_1^\perp \cup \dots \cup H_n^\perp)$.*

Note that if $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ is $\text{GL}(n)$ covariant (contravariant), then the extended operator Z is $\text{GL}(n)$ covariant (contravariant) on $\overline{\mathcal{P}}_0$.

2.2 Covariant valuations

We prove the following result.

Proposition 1. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ covariant of weight $q \geq 0$. Then there are constants $c_1, c_2 \geq 0$ such that*

$$Z P = c_1 P \tilde{+} c_2(-P)$$

for every $P \in \mathcal{P}_0$.

To extend Z to $\overline{\mathcal{P}}_0$, we apply Lemmas 1 and 2 and need the following result.

Lemma 5. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ covariant of weight q and let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be defined by $Y P(\cdot) = \rho(Z P, \cdot)$. Then for every central hyperplane H and $P \in \mathcal{P}_0(H)$, the following holds: If $q > -1$, then Y is vanishing on $S^{n-1} \setminus H$ at P . If $q > 0$, then Y is vanishing on S^{n-1} at P . If $q \geq 0$, then Y is bounded at P .*

Proof. Since Z is rotation covariant, it suffices to prove the statements for $H = e_n^\perp$. Let $P \in \mathcal{P}_0(H)$. Let $u \in H^- \setminus H$ and $v \in H^+ \setminus H$ be chosen such that $[P, u, v] = [P, u] \cup [P, v]$ and let $r > 0$ be suitably small. Since Z is a valuation, we have

$$Z[P, u, v] + Z[P, -r u, -r v] = Z[P, u, -r u] + Z[P, v, -r v]$$

and

$$Z[P, u, -r u] + Z[P, r u, -u] = Z[P, u, -u] + Z[P, r u, -r u].$$

Thus to prove the lemma it suffices to show that $Y[P, u, -u]$ is bounded on S^{n-1} for $|u| \leq 1$ for $q \geq 0$ and that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ locally uniformly on $S^{n-1} \setminus H$ for $q > -1$ or uniformly on S^{n-1} for $q > 0$.

Define $\phi_u \in \text{GL}(n)$ by $\phi_u e_j = e_j$, $j = 1, \dots, n-1$, and $\phi_u e_n = u$. Then

$$\phi_u^{-1}x = \left(x_1 - \frac{u_1}{u_n}x_n, \dots, x_{n-1} - \frac{u_{n-1}}{u_n}x_n, \frac{x_n}{u_n}\right). \quad (22)$$

Since Z is $\text{GL}(n)$ covariant of weight q , we obtain by (1)

$$Z[P, u, -u](x) = Z(\phi_u[P, e_n, -e_n])(x) = u_n^q Z[P, e_n, -e_n](\phi_u^{-1}x).$$

and

$$Y[P, u, -u](x) = u_n^{q+1} Y[P, e_n, -e_n](u_n x_1 - u_1 x_n, \dots, u_n x_{n-1} - u_{n-1} x_n, x_n). \quad (23)$$

For $q > -1$, this implies that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ locally uniformly on $S^{n-1} \setminus H$. If $x \in S^{n-1} \cap H$, it follows from (23) and (1) that

$$Y[P, u, -u](x) = u_n^q Y[P, e_n, -e_n](x_1, \dots, x_{n-1}, 0).$$

Thus, we obtain that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ uniformly on S^{n-1} for $q > 0$. Let $x \in S^{n-1}$. It follows from (23) and (1) that

$$Y[P, u, -u](x) \leq \frac{u_n^{q+1}}{|\phi_u^{-1}x|} \max_{w \in S^{n-1}} Y[P, e_n, -e_n](w). \quad (24)$$

If $|u_n| \geq 4|x_n|$, then by (22)

$$|\phi_u^{-1}x| = \left|x + \frac{x_n}{u_n}(e_n - u)\right| \geq 1 - \frac{1}{4}|e_n - u| \geq \frac{1}{2}.$$

If $|u_n| \leq 4|x_n|$, then by (22), $|\phi_u^{-1}x| \geq |(\phi_u^{-1}x) \cdot e_n| \geq \frac{1}{4}$. Thus (24) implies that Y is bounded at P for $q \geq 0$. \square

We also write Z for the extended operator. Let T be the simplex with vertices $0, e_1, \dots, e_n$. We determine ZT . Since Z is $\text{GL}(n)$ covariant,

$$\rho(ZT, (x_1, \dots, x_n)) = \rho(ZT, (x_{i_1}, \dots, x_{i_n})) \quad (25)$$

for every permutation (i_1, \dots, i_n) of $(1, \dots, n)$. Let Z' be a simple and $\text{GL}(n)$ covariant valuation on $\overline{\mathcal{P}}_0$. Note that it suffices to show that $ZT = Z'T$ to show that $ZP = Z'P$ for every $P \in \mathcal{P}_0$. This implies that Proposition 1 is a consequence of the following lemmas.

First, let $q > 0$. Note that in this case it follows from Lemma 2 and Lemma 5 that $\rho(ZT, \cdot)$ is continuous on S^{n-1} .

Lemma 6. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ covariant of weight $q > 0$. Then $ZT = \{0\}$.*

Proof. For $0 < \lambda_j < 1$, $j = 2, \dots, n$, let H_j be the central hyperplane with normal vector $\lambda_j e_1 - (1 - \lambda_j) e_j$. Then H_j dissects T into two simplices $T \cap H_j^+$ and $T \cap H_j^-$. Since Z is a simple valuation, we have

$$ZT = Z(T \cap H_j^+) \dot{+} Z(T \cap H_j^-). \quad (26)$$

For $j = 2, \dots, n$, define ϕ_j, ψ_j by

$$\begin{aligned}\phi_j e_j &= (1 - \lambda_j) e_1 + \lambda_j e_j & \text{and} & & \phi_j e_i &= e_i & \text{for } i \neq j, \\ \psi_j e_1 &= (1 - \lambda_j) e_1 + \lambda_j e_j & \text{and} & & \psi_j e_i &= e_i & \text{for } i \neq 1.\end{aligned}$$

Then $T \cap H_j^+ = \phi_j T$ and $T \cap H_j^- = \psi_j T$. Set $f(x) = \rho(ZT, x)$ and let $x \neq 0$. Since Z is $\text{GL}(n)$ covariant of weight q , (26) and (1) imply that

$$f(x) = \lambda_j^q f(\phi_j^{-1} x) + (1 - \lambda_j)^q f(\psi_j^{-1} x). \quad (27)$$

Note that

$$\begin{aligned}\phi_j^{-1} e_j &= -\frac{1 - \lambda_j}{\lambda_j} e_1 + \frac{1}{\lambda_j} e_j & \text{and} & & \phi_j^{-1} e_i &= e_i & \text{for } i \neq j, \\ \psi_j^{-1} e_1 &= \frac{1}{1 - \lambda_j} e_1 - \frac{\lambda_j}{1 - \lambda_j} e_j & \text{and} & & \psi_j^{-1} e_i &= e_i & \text{for } i \neq 1.\end{aligned}$$

From (27) with $x = e_1$, we obtain

$$f(e_1 - \lambda_j e_j) = \frac{1 - \lambda_j^q}{(1 - \lambda_j)^{q+1}} f(e_1). \quad (28)$$

Since $(1 - \lambda_j^q)/(1 - \lambda_j)^{q+1} \rightarrow \infty$ as $\lambda_j \rightarrow 1$, we obtain that $f(e_1) = 0$ and by (25) that $f(e_i) = 0$, $i = 1, \dots, n$. Similarly, we obtain that $f(-e_i) = 0$, $i = 1, \dots, n$. From (27), we obtain

$$f((1 - \lambda_j) e_1 + \lambda_j e_j) = \lambda_j^q f(e_j) + (1 - \lambda_j)^q f(e_1).$$

It follows from this and (28) that $f(x_1 e_1 + x_j e_j) = 0$ for every $x_1, x_j \in \mathbb{R}$, $(x_1, x_j) \neq (0, 0)$. Let $x' = x_2 e_2 + \dots + x_{j-1} e_{j-1}$. Then by (27)

$$f((1 - \lambda_j) e_1 + \lambda_j e_j + x') = \lambda_j^q f(e_j + x') + (1 - \lambda_j)^q f(e_1 + x'). \quad (29)$$

By (25), $f(e_1 + x') = f(e_j + x')$. Thus by using induction on the number of vanishing coordinates, we obtain from (29) that $f(x) = 0$ for every $x \neq 0$. \square

Next, we consider the case $q = 0$. Note that in this case it follows from Lemma 1 and Lemma 5 that $\rho(ZT, \cdot)$ is continuous and uniformly bounded on $S^{n-1} \setminus (e_1^\perp \cup \dots \cup e_n^\perp)$.

Lemma 7. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ covariant of weight $q = 0$. Then there are constants $c_1, c_2 \geq 0$ such that*

$$\rho(ZT, x) = c_1 \rho(T, x) + c_2 \rho(-T, x)$$

for $x \notin e_1^\perp \cup \dots \cup e_n^\perp$.

Proof. We define H_j, ϕ_j, ψ_j , and f as in the proof of Lemma 6. Let $x \notin e_1^\perp \cup \dots \cup e_n^\perp$. Note that for given x , there is a dense set of λ_2 such that the subsequent expressions are well defined, that is, for example, $\phi_2^{-1} x \notin e_1^\perp \cup \dots \cup e_n^\perp$. As in (27), we have

$$f(x) = f(\phi_2^{-1} x) + f(\psi_2^{-1} x). \quad (30)$$

Using this repeatedly, we obtain

$$f(\psi_2^k x) = \sum_{j=1}^k f(\phi_2^{-1} \psi_2^j x) + f(x). \quad (31)$$

Let $x' = x_2 e_2 + \cdots + x_n e_n$. Note that for ψ_2 the vectors e_i , $i = 2, \dots, n$, are eigenvectors with eigenvalue 1 and the vector $e_1 - e_2$ is an eigenvector with eigenvalue $(1 - \lambda_2)$. Setting $x = (e_1 - e_2) + x'$ in (31) gives

$$f((1 - \lambda_2)^k (e_1 - e_2) + x') = \sum_{j=1}^k f(\phi_2^{-1} ((1 - \lambda_2)^j (e_1 - e_2) + x')) + f(x).$$

Let $k \rightarrow \infty$. Since f is uniformly bounded, continuous and non-negative, we obtain that $f(\phi_2^{-1} x') = 0$. Note that

$$\phi_2^{-1} x' = -\frac{1 - \lambda_2}{\lambda_2} x_2 e_1 + \frac{1}{\lambda_2} x_2 e_2 + x_3 e_3 + \cdots + x_n e_n.$$

Using (25) and the continuity of f , we conclude that

$$f(x_1, \dots, x_n) = 0 \text{ if } x_i \neq 0 \text{ for } i = 1, \dots, n, \text{ and not all } x_i \text{ have the same sign.} \quad (32)$$

Let $x_1, \dots, x_n > 0$. Then by (30) and (32) we have

$$f(\phi_2 x) = f(x) + f(\psi_2^{-1} \phi_2 x) = f(x).$$

Thus

$$f(\phi_n \cdots \phi_2 x) = f(\phi_{n-1} \cdots \phi_2 x) = \cdots = f(x).$$

Since

$$\phi_n \cdots \phi_2 (x_1, \dots, x_n) = (x_1 + (1 - \lambda_2) x_2 + \cdots + (1 - \lambda_n) x_n, \lambda_2 x_2, \dots, \lambda_n x_n), \quad (33)$$

we obtain that

$$f(x_1, \dots, x_n) = f(1, \dots, 1) \text{ for } x_1 + \cdots + x_n = n, 0 < x_2, \dots, x_n < 1.$$

By choosing λ_i such that $\lambda_i x_i < 1$, we obtain from this and (33) that

$$f(x_1, \dots, x_n) = f(1, \dots, 1) \text{ for } x_1 + \cdots + x_n = n, x_1, \dots, x_n > 0.$$

Similarly, we obtain that

$$f(-x_1, \dots, -x_n) = f(-1, \dots, -1) \text{ for } x_1 + \cdots + x_n = n, x_1, \dots, x_n > 0.$$

Thus $f(x) = c_1 \rho(T, x) + c_2 \rho(-T, x)$ with suitable constants $c_1, c_2 \geq 0$. \square

2.3 Contravariant valuations

We prove the following result.

Proposition 2. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\mathrm{GL}(n)$ contravariant of weight $q > 0$. Then there is a constant $c \geq 0$ such that*

$$ZP = cIP$$

for every $P \in \mathcal{P}_0$.

To extend Z to $\overline{\mathcal{P}_0}$, we apply Lemmas 2, 3 and 4 and need the following result.

Lemma 8. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\mathrm{GL}(n)$ contravariant of weight q and let $Y : \mathcal{P}_0 \rightarrow \mathcal{C}_+(S^{n-1})$ be defined by $YP(\cdot) = \rho(ZP, \cdot)$. For every central hyperplane H , the following holds: If $q > 0$, then Y is vanishing on $S^{n-1} \setminus H^\perp$ at $\mathcal{P}_0(H)$. If $q > 1$, then Y is vanishing on S^{n-1} at $\mathcal{P}_0(H)$. If $q \geq 1$, then Y is bounded on S^{n-1} at $\mathcal{P}_0(H)$.*

Proof. Since Z is rotation contravariant, it suffices to prove the statements for $H = e_n^\perp$. Let $P \in \mathcal{P}_0(H)$. Let $u \in H^- \setminus H$, $v \in H^+ \setminus H$ be chosen such that $[P, u, v] = [P, u] \cup [P, v]$ and let $r > 0$ be suitably small. Since Z is a valuation, we have

$$Z[P, u, v] + Z[P, -ru, -rv] = Z[P, u, -ru] + Z[P, v, -rv]$$

and

$$Z[P, u, -ru] + Z[P, ru, -u] = Z[P, u, -u] + Z[P, ru, -ru].$$

Thus to prove the lemma it suffices to show that $Y[P, u, -u]$ is bounded on S^{n-1} for $|u| \leq 1$ for $q \geq 1$ and that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ locally uniformly on $S^{n-1} \setminus H^\perp$ for $q > 0$ or uniformly on S^{n-1} for $q > 1$.

Define $\phi_u \in \mathrm{GL}(n)$ by $\phi_u e_j = e_j$, $j = 1, \dots, n-1$, and $\phi_u e_n = u$. Then

$$\phi_u^t x = (x_1, \dots, x_{n-1}, x_1 u_1 + \dots + x_n u_n). \quad (34)$$

Since Z is $\mathrm{GL}(n)$ contravariant of weight q , we obtain by (1)

$$Z[P, u, -u](x) = Z(\phi_u[P, e_n, -e_n])(x) = u_n^q Z[P, e_n, -e_n](\phi_u^t x).$$

and

$$Y[P, u, -u](x) = u_n^q Y[P, e_n, -e_n](x_1, \dots, x_{n-1}, x_1 u_1 + \dots + x_n u_n). \quad (35)$$

For $q > 0$, this implies that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ locally uniformly on $S^{n-1} \setminus H^\perp$. If $x \in S^{n-1} \cap H^\perp$, it follows from (35) and (1) that

$$Y[P, u, -u](x) = u_n^{q-1} Y[P, e_n, -e_n](0, \dots, 0, x_n).$$

Thus, we obtain that $\lim_{u \rightarrow 0} Y[P, u, -u] = 0$ uniformly on S^{n-1} for $q > 1$. Let $x \in S^{n-1}$. It follows from (35) and (1) that

$$Y[P, u, -u](x) \leq \frac{u_n^q}{|\phi_u^t x|} \max_{w \in S^{n-1}} Y[P, e_n, -e_n](w). \quad (36)$$

Since $|u_n| \leq 1$, by (34)

$$\frac{|\phi_u^t x|^2}{u_n^2} = \frac{x_1^2}{u_n^2} + \dots + \frac{x_{n-1}^2}{u_n^2} + \frac{(x \cdot u)^2}{u_n^2} \geq 1.$$

Thus (36) implies that Y is bounded at P for $q \geq 0$. \square

We also write Z for the extended operator. Let T be the simplex with vertices $0, e_1, \dots, e_n$. We determine ZT . Since Z is $\text{GL}(n)$ contravariant,

$$\rho(ZT, (x_1, \dots, x_n)) = \rho(ZT, (x_{i_1}, \dots, x_{i_n})) \quad (37)$$

for every permutation (i_1, \dots, i_n) of $(1, \dots, n)$. Let Z' be a simple and $\text{GL}(n)$ contravariant valuation on $\overline{\mathcal{P}}_0$. Note that it suffices to show that $ZT = Z'T$ to show that $ZP = Z'P$ for every $P \in \mathcal{P}_0$. This implies that Proposition 2 is a consequence of the following lemmas.

First, let $q > 1$. Note that in this case it follows from Lemma 2 and Lemma 8 that $\rho(ZT, \cdot)$ is continuous on S^{n-1} .

Lemma 9. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ contravariant of weight $q > 1$. Then $ZT = \{0\}$.*

Proof. We define H_j, ϕ_j, ψ_j , and f as in the proof of Lemma 6. Let $x \neq 0$. Since Z is $\text{GL}(n)$ contravariant of weight q , (26) and (1) imply that

$$f(x) = \lambda_j^q f(\phi_j^t x) + (1 - \lambda_j)^q f(\psi_j^t x). \quad (38)$$

From this with $x = e_1$, we obtain

$$f(e_1 + (1 - \lambda_j)e_j) = \frac{1 - (1 - \lambda_j)^{q-1}}{\lambda_j^q} f(e_1). \quad (39)$$

Since $(1 - (1 - \lambda_j)^{q-1})/\lambda_j^q \rightarrow \infty$ as $\lambda_j \rightarrow 0$, we obtain that $f(e_1) = 0$ and by (37) that $f(e_i) = 0$, $i = 1, \dots, n$. Similarly, we obtain that $f(-e_i) = 0$, $i = 1, \dots, n$. From (38), we obtain

$$f(\lambda_j e_1 - (1 - \lambda_j)e_j) = \lambda_j^{q-1} f(e_1) + (1 - \lambda_j)^{q-1} f(-e_j).$$

It follows from this and (39) that $f(x_1 e_1 + x_j e_j) = 0$ for every $x_1, x_j \in \mathbb{R}$, $(x_1, x_j) \neq (0, 0)$. Let $x' = x_2 e_2 + \dots + x_{j-1} e_{j-1}$. Then by (38)

$$f(\lambda_j e_1 - (1 - \lambda_j)e_j + x') = \lambda_j^q f(\lambda_j e_1 + x') + (1 - \lambda_j)^q f(-(1 - \lambda_j)e_j + x'). \quad (40)$$

By (37), $f(-(1 - \lambda_j)e_j + x') = f(-(1 - \lambda_j)e_1 + x')$. Thus by using induction on the number of vanishing coordinates, we obtain from (40) that $f(x) = 0$ for every $x \neq 0$. \square

Next, we consider the case $0 < q < 1$. Let \mathbb{R}_a^n be the set of $x \in \mathbb{R}^n$ not on the coordinate axes, that is, $x \in \mathbb{R}^n$, $x \neq \lambda e_j$, $\lambda \in \mathbb{R}$, $j = 1, \dots, n$. Note that for $0 < q < 1$ it follows from Lemma 4 and Lemma 8 that $\rho(ZT, \cdot)$ is continuous on \mathbb{R}_a^n .

Lemma 10. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ contravariant of weight $0 < q < 1$. Then*

$$\rho(ZT, x) = 0$$

for $x \in \mathbb{R}_a^n$.

Proof. We define H_j, ϕ_j, ψ_j , and f as in the proof of Lemma 6. We consider $x \in \mathbb{R}_a^n$ for which the subsequent expressions are well defined, that is, for example, $\phi_2^t x \in \mathbb{R}_a^n$. Since Z is $\text{GL}(n)$ contravariant of weight q , (26) and (1) imply that

$$f(x) = \lambda_2^q f(\phi_2^t x) + (1 - \lambda_2)^q f(\psi_2^t x). \quad (41)$$

For $x_2 = 1 - \lambda_2$, it follows that

$$\begin{aligned} f(x_1, 1 - \lambda_2, x_3, \dots, x_n) &= \lambda_2^q f(x_1, (x_1 + \lambda_2)(1 - \lambda_2), x_3, \dots, x_n) \\ &\quad + (1 - \lambda_2)^q f((x_1 + \lambda_2)(1 - \lambda_2), (1 - \lambda_2), x_3, \dots, x_n). \end{aligned}$$

Let $x_1 \rightarrow -\lambda_2, x_3, \dots, x_n \rightarrow 0$. Since the left hand side is well defined and f is non-negative, this implies that f is uniformly bounded on $S^{n-1} \cap \mathbb{R}_a^n$. By (41),

$$f(\phi_2^{-t}x) = \lambda_2^q f(x) + (1 - \lambda_2)^q f(\psi_2^t \phi_2^{-t}x).$$

Setting $x_1 = -1, x_2 = 1$ and dividing by λ_2 gives

$$\begin{aligned} f(-\lambda_2, 2 - \lambda_2, \lambda_2 x_3, \dots, \lambda_2 x_n) \\ = \lambda_2^{q-1} f(-1, 1, x_3, \dots, x_n) + (1 - \lambda_2)^q f(\lambda_2, 2 - \lambda_2, \lambda_2 x_3, \dots, \lambda_2 x_n). \end{aligned}$$

Let $\lambda_2 \rightarrow 0$. Since f is uniformly bounded on $S^{n-1} \cap \mathbb{R}_a^n$, this shows that

$$f(-1, 1, x_3, \dots, x_n) = 0.$$

Combined with (41) for $x = (-1, 1, x_3, \dots, x_n)$ this gives

$$0 = \lambda_2^q f(-1, -1 + 2\lambda_2, x_3, \dots, x_n) + (1 - \lambda_2)^q f(-1 + 2\lambda_2, 1, x_3, \dots, x_n).$$

Since f is non-negative, this implies that

$$f(-1, -1 + 2\lambda_2, x_3, \dots, x_n) = f(-1 + 2\lambda_2, 1, x_3, \dots, x_n) = 0.$$

Similarly, we obtain that $f(1, -1, x_3, \dots, x_n) = 0$ and conclude that $f(x) = 0$ for every $x \in \mathbb{R}_a^n$. \square

Finally, we consider the case $q = 1$. Note that in this case it follows from Lemma 3 and Lemma 8 that $\rho(ZT, \cdot)$ is continuous and uniformly bounded on $S^{n-1} \cap \mathbb{R}_a^n$.

Lemma 11. *Let $Z : \mathcal{P}_0 \rightarrow \mathcal{S}$ be a valuation which is $\text{GL}(n)$ contravariant of weight $q = 1$. Then there is a constant $c \geq 0$ such that*

$$\rho(ZT, x) = c \rho(IT, x)$$

for $x \in \mathbb{R}_a^n$.

Proof. We define H_j, ϕ_j, ψ_j , and f as in the proof of Lemma 6. Let $x \in \mathbb{R}_a^n$. Note that for given x , there is a dense set of λ_j such that the subsequent expressions are well defined, that is, for example, $\phi_j^t x \in \mathbb{R}_a^n$. Since Z is $\text{GL}(n)$ contravariant of weight $q = 1$, (26) and (1) imply that

$$f(x) = \lambda_j f(\phi_j^t x) + (1 - \lambda_j) f(\psi_j^t x). \quad (42)$$

Using this repeatedly, we obtain

$$f(x) = \lambda_2 \cdots \lambda_n f(\phi_n^t \cdots \phi_2^t x) + \sum_{j=3}^n \lambda_2 \cdots \lambda_{j-1} (1 - \lambda_j) f(\psi_j^t \phi_{j-1}^t \cdots \phi_2^t x) + (1 - \lambda_2) f(\psi_2^t x).$$

Using this repeatedly, we obtain

$$\begin{aligned}
f((\psi_2^{-t})^k x) &= \lambda_2 \cdots \lambda_n \sum_{i=1}^k (1 - \lambda_j)^{k-i} f(\phi_n^t \cdots \phi_2^t (\psi_2^{-t})^i x) \\
&+ \sum_{j=3}^n \lambda_2 \cdots \lambda_{j-1} \sum_{i=1}^k (1 - \lambda_j)^{k-i} f(\psi_j^t \phi_{j-1}^t \cdots \phi_2^t (\psi_2^{-t})^i x) + (1 - \lambda_2)^k f(x).
\end{aligned} \tag{43}$$

Let $x' = x_3 e_3 + \cdots + x_n e_n$. Note that for ψ_2^t the vectors $e_1 + e_2$ and e_i , $i = 3, \dots, n$, are eigenvectors with eigenvalue 1 and the vector e_1 is an eigenvector with eigenvalue $(1 - \lambda_2)$. Setting $x = e_1 + x'$ in (43) and dividing by $(1 - \lambda_2)^k$ gives

$$\begin{aligned}
f(e_1 + (1 - \lambda_2)^k x') &= \lambda_2 \cdots \lambda_n \sum_{i=1}^k f(\phi_n^t \cdots \phi_2^t (e_1 + (1 - \lambda_2)^i x')) \\
&+ \sum_{j=3}^n \lambda_2 \cdots \lambda_{j-1} \sum_{i=1}^k f(\psi_j^t \phi_{j-1}^t \cdots \phi_2^t (e_1 + (1 - \lambda_2)^i x')) + f(x).
\end{aligned}$$

Let $k \rightarrow \infty$. Since f is uniformly bounded, continuous and non-negative, we obtain that $f(\phi_n^t \cdots \phi_2^t e_1) = 0$. Note that $\phi_n^t \cdots \phi_2^t e_1 = e_1 + (1 - \lambda_2) e_2 + \cdots + (1 - \lambda_n) e_n$. Thus we conclude that

$$f(x_1, \dots, x_n) = 0 \quad \text{for } x_1, \dots, x_n > 0. \tag{44}$$

Next, let $x_1 < 0, x_2, \dots, x_n > 0$ and $(1 - \lambda_j) x_1 + \lambda_j x_j > 0$, $j = 2, \dots, n$. Then by (42) and (44)

$$f(x) = \lambda_2 f(\phi_2^t x) + (1 - \lambda_2) f(\psi_2^t x) = \lambda_2 f(\phi_2^t x)$$

and

$$f(\phi_n^t \cdots \phi_2^t x) = \frac{1}{\lambda_2 \cdots \lambda_n} f(x).$$

Setting $x = (-1, 1, \dots, 1)$ and $a = f(-1, 1, \dots, 1)$, we obtain

$$f(-1, -1 + 2\lambda_2, \dots, -1 + 2\lambda_n) = \frac{a}{\lambda_2 \cdots \lambda_n}.$$

We use induction on the number of negative coordinates. Suppose f is determined by a for x with at most $(k-1)$ negative coordinates. Let $x_1, x_2 < 0$ and $(1 - \lambda_2) x_1 + \lambda_2 x_2 > 0$. Then

$$f(x) = \lambda_2 f(\phi_2^t x) + (1 - \lambda_2) f(\psi_2^t x)$$

and $\phi_2^t x$ as well as $\psi_2^t x$ have at most $(k-1)$ negative coordinates. Thus $f(x)$ is determined by a . We obtain that for given a there is at most one operator Z . Since $Z = c I$ with a suitable $c \geq 0$ is such an operator, this concludes the proof of the lemma. \square

Acknowledgements

The author is grateful to the referees for the careful reading given to the original draft of this paper and for the suggested improvements. The author also thanks Erwin Lutwak and Vitali Milman for their help with the presentation of this result.

References

- [1] S. Alesker, *Continuous rotation invariant valuations on convex sets*, Ann. of Math. (2) **149** (1999), 977–1005.
- [2] S. Alesker, *Description of translation invariant valuations on convex sets with solution of P. McMullen’s conjecture*, Geom. Funct. Anal. **11** (2001), 244–272.
- [3] S. Alesker, *Hard Lefschetz theorem for valuations, complex integral geometry, and unitarily invariant valuations*, J. Differential Geom. **63** (2003), 63–95.
- [4] S. Alesker, *The multiplicative structure on continuous polynomial valuations*, Geom. Funct. Anal. **14** (2004), 1–26.
- [5] H. Busemann, *A theorem on convex bodies of the Brunn-Minkowski type*, Proc. Nat. Acad. Sci. U. S. A. **35** (1949), 27–31.
- [6] H. Busemann, *Volume in terms of concurrent cross-sections*, Pacific J. Math. **3** (1953), 1–12.
- [7] R. Gardner, *Intersection bodies and the Busemann-Petty problem*, Trans. Amer. Math. Soc. **342** (1994), 435–445.
- [8] R. Gardner, *A positive answer to the Busemann-Petty problem in three dimensions*, Ann. of Math. (2) **140** (1994), 435–447.
- [9] R. Gardner, *Geometric tomography*, Cambridge University Press, Cambridge, 1995.
- [10] R. Gardner, A. Koldobsky, and T. Schlumprecht, *An analytic solution to the Busemann-Petty problem on sections of convex bodies*, Ann. of Math. (2) **149** (1999), 691–703.
- [11] P. Goodey, E. Lutwak, and W. Weil, *Functional analytic characterizations of classes of convex bodies*, Math. Z. **222** (1996), 363–381.
- [12] E. Grinberg and G. Zhang, *Convolutions, transforms, and convex bodies*, Proc. London Math. Soc. (3) **78** (1999), 77–115.
- [13] H. Hadwiger, *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*, Springer, Berlin, 1957.
- [14] N. J. Kalton and A. Koldobsky, *Intersection bodies and L_p spaces*, Adv. Math. **196** (2005), 257–275.
- [15] D. A. Klain, *Star valuations and dual mixed volumes*, Adv. Math. **121** (1996), 80–101.
- [16] D. A. Klain, *Invariant valuations on star-shaped sets*, Adv. Math. **125** (1997), 95–113.
- [17] D. A. Klain and G.-C. Rota, *Introduction to geometric probability*, Cambridge University Press, Cambridge, 1997.
- [18] A. Koldobsky, *Intersection bodies in \mathbf{R}^4* , Adv. Math. **136** (1998), 1–14.
- [19] A. Koldobsky, *Intersection bodies, positive definite distributions, and the Busemann-Petty problem*, Amer. J. Math. **120** (1998), 827–840.
- [20] A. Koldobsky, *Second derivative test for intersection bodies*, Adv. Math. **136** (1998), 15–25.
- [21] A. Koldobsky, *A functional analytic approach to intersection bodies*, Geom. Funct. Anal. **10** (2000), 1507–1526.
- [22] A. Koldobsky, *Fourier analysis in convex geometry*, Mathematical Surveys and Monographs, vol. 116, American Mathematical Society, Providence, RI, 2005.
- [23] K. Leichtweiß, *Affine geometry of convex bodies*, Johann Ambrosius Barth, Heidelberg, 1998.
- [24] M. Ludwig, *Projection bodies and valuations*, Adv. Math. **172** (2002), 158–168.
- [25] M. Ludwig, *Valuations on polytopes containing the origin in their interiors*, Adv. Math. **170** (2002), 239–256.
- [26] M. Ludwig, *Ellipsoids and matrix valued valuations*, Duke Math. J. **119** (2003), 159–188.
- [27] M. Ludwig, *Minkowski valuations*, Trans. Amer. Math. Soc. **357** (2005), 4191–4213.
- [28] M. Ludwig and M. Reitzner, *A characterization of affine surface area*, Adv. Math. **147** (1999), 138–172.
- [29] E. Lutwak, *Intersection bodies and dual mixed volumes*, Adv. Math. **71** (1988), 232–261.

- [30] E. Lutwak, *On some ellipsoid formulas of Busemann, Furstenberg and Tzkonik, Guggenheimer, and Petty*, J. Math. Anal. Appl. **159** (1991), 18–26.
- [31] E. Lutwak, *Selected affine isoperimetric inequalities*, Handbook of Convex Geometry, Vol. A (P.M. Gruber and J.M. Wills, eds.), North-Holland, Amsterdam, 1993, 151–176.
- [32] P. McMullen, *Valuations and dissections*, Handbook of Convex Geometry, Vol. B (P.M. Gruber and J.M. Wills, eds.), North-Holland, Amsterdam, 1993, 933–990.
- [33] P. McMullen and R. Schneider, *Valuations on convex bodies*, Convexity and its applications (P.M. Gruber and J.M. Wills, eds.), Birkhäuser, 1983, 170–247.
- [34] V. D. Milman and A. Pajor, *Isotropic position and inertia ellipsoids and zonoids of the unit ball of a normed n -dimensional space*, Geometric aspects of functional analysis (1987–88), Lecture Notes in Math., vol. 1376, Springer, Berlin, 1989, 64–104.
- [35] B. Rubin and G. Zhang, *Generalizations of the Busemann-Petty problem for sections of convex bodies*, J. Funct. Anal. **213** (2004), 473–501.
- [36] R. Schneider, *Convex bodies: the Brunn–Minkowski theory*, Cambridge Univ. Press, Cambridge, 1993.
- [37] A. C. Thompson, *Minkowski geometry*, Cambridge University Press, Cambridge, 1996.
- [38] G. Zhang, *Centered bodies and dual mixed volumes*, Trans. Am. Math. Soc. **345** (1994), 777–801.
- [39] G. Zhang, *A positive solution to the Busemann-Petty problem in \mathbf{R}^4* , Ann. of Math. (2) **149** (1999), 535–543.

Institut für Diskrete Mathematik und Geometrie
 Technische Universität Wien
 Wiedner Hauptstraße 8-10/104
 1040 Wien, Austria
 E-mail: monika.ludwig@tuwien.ac.at