# Möbius Orthogonality for the Zeckendorf Sum-of-Digits Function

#### Clemens Müllner







20. December 2018

Joint work with Michael Drmota and Lukas Spiegelhofer



#### Möbius function

The Möbius function is defined by

$$\mu(n) = \left\{ egin{array}{ll} (-1)^k & ext{if $n$ is squarefree and} \\ k & ext{is the number of prime factors} \\ 0 & ext{otherwise} \end{array} 
ight.$$

A sequence **u** is **orthogonal to the Möbius function**  $\mu$ (n) if

$$\sum_{n\leq N}\mu(n)u_n=o(\sum_{n\leq N}|u_n|)\qquad (N\to\infty).$$

#### Old Heuristic - Mobius Randomness Law

Any "reasonably defined (easy)"bounded sequence independent of  $\mu$  is orthogonal to  $\mu$ .



#### Möbius function

The Möbius function is defined by

$$\mu(n) = \left\{ egin{array}{ll} (-1)^k & ext{if $n$ is squarefree and} \\ k & ext{is the number of prime factors} \\ 0 & ext{otherwise} \end{array} 
ight.$$

A sequence  $\mathbf{u}$  is orthogonal to the Möbius function  $\mu(n)$  if

$$\sum_{n\leq N}\mu(n)u_n=o(\sum_{n\leq N}|u_n|)\qquad (N\to\infty).$$

#### Old Heuristic - Mobius Randomness Law

Any "reasonably defined (easy)"bounded sequence independent of  $\mu$  is orthogonal to  $\mu$ .



#### Möbius function

The Möbius function is defined by

$$\mu(n) = \left\{ egin{array}{ll} (-1)^k & ext{if $n$ is squarefree and} \\ k & ext{is the number of prime factors} \\ 0 & ext{otherwise} \end{array} 
ight.$$

A sequence **u** is **orthogonal to the Möbius function**  $\mu$ (n) if

$$\sum_{n\leq N}\mu(n)u_n=o(\sum_{n\leq N}|u_n|)\qquad (N\to\infty).$$

#### Old Heuristic - Mobius Randomness Law

Any "reasonably defined (easy)"bounded sequence independent of  $\mu$  is orthogonal to  $\mu.$ 

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

- Constant sequences ⇔ PNT
- Periodic sequences ⇔ PNT in arithmetic Progressions
- Quasiperiodic sequences  $f(n) = F(\alpha n \mod 1)$  Davenport
- Nilsequences Green and Tao
- Horocycle Flows Bourgain, Sarnak and Ziegler
- Dynamical systems with discrete spectrum

### Sarnak Conjecture

#### Definition

A dynamical system is said to be deterministic, if its topological entropy is 0.

#### Conjecture (Sarnak conjecture, 2010)

Every bounded complex sequence  $\mathbf{u} = (u_n)_{n>0}$  that is obtained by a deterministic dynamical system is orthogonal to the Möbius function  $\mu(n)$ .

### Sarnak Conjecture

#### Definition

A dynamical system is said to be deterministic, if its topological entropy is 0.

#### Conjecture (Sarnak conjecture, 2010)

Every bounded complex sequence  $\mathbf{u} = (u_n)_{n>0}$  that is obtained by a deterministic dynamical system is orthogonal to the Möbius function  $\mu(n)$ .

### Chowla Conjecture

#### Conjecture (Chowla)

Let  $0 \le a_1 < a_2 < \ldots < a_t$  and  $k_1, k_2, \ldots, k_t$  in  $\{1,2\}$  not all even, then as  $N \to \infty$ 

$$\sum_{n\leq N} \mu^{k_1}(n+a_1)\mu^{k_2}(n+a_2)\cdots \mu^{k_t}(n+a_t) = o(N).$$

#### Theorem (Sarnak)

The Chowla Conjecture implies the Sarnak Conjecture.

#### Theorem (Tao)

The logarithmic version of the Sarnak Conjecture implies the logarithmic version of the Chowla Conjecture.



### Chowla Conjecture

#### Conjecture (Chowla)

Let  $0 \le a_1 < a_2 < \ldots < a_t$  and  $k_1, k_2, \ldots, k_t$  in  $\{1,2\}$  not all even, then as  $N \to \infty$ 

$$\sum_{n\leq N} \mu^{k_1}(n+a_1)\mu^{k_2}(n+a_2)\cdots \mu^{k_t}(n+a_t) = o(N).$$

#### Theorem (Sarnak)

The Chowla Conjecture implies the Sarnak Conjecture.

#### Theorem (Tao

The logarithmic version of the Sarnak Conjecture implies the logarithmic version of the Chowla Conjecture.

### Chowla Conjecture

#### Conjecture (Chowla)

Let  $0 \le a_1 < a_2 < \ldots < a_t$  and  $k_1, k_2, \ldots, k_t$  in  $\{1,2\}$  not all even, then as  $N \to \infty$ 

$$\sum_{n\leq N} \mu^{k_1}(n+a_1)\mu^{k_2}(n+a_2)\cdots \mu^{k_t}(n+a_t) = o(N).$$

#### Theorem (Sarnak)

The Chowla Conjecture implies the Sarnak Conjecture.

#### Theorem (Tao)

The logarithmic version of the Sarnak Conjecture implies the logarithmic version of the Chowla Conjecture.



### Logarithmic versions

#### Theorem (Tao, Tao - Teräväinen)

The logarithmic version of the Chowla conjecture is true for t=2 and for t odd.

#### Theorem (Frantzikinakis, Host)

The logarithmic version of the Sarnak conjecture is true if the dynamical system has countable many ergodic components.

Literature: "Sarnak conjecture: What's new?" (Ferenczi - Kulaga Przymus - Lemanczyk)

### Logarithmic versions

#### Theorem (Tao, Tao - Teräväinen)

The logarithmic version of the Chowla conjecture is true for t=2 and for t odd.

#### Theorem (Frantzikinakis, Host)

The logarithmic version of the Sarnak conjecture is true if the dynamical system has countable many ergodic components.

Literature: "Sarnak conjecture: What's new?" (Ferenczi - Kulaga Przymus - Lemanczyk)

### Logarithmic versions

#### Theorem (Tao, Tao - Teräväinen)

The logarithmic version of the Chowla conjecture is true for t=2 and for t odd.


#### Theorem (Frantzikinakis, Host)

The logarithmic version of the Sarnak conjecture is true if the dynamical system has countable many ergodic components.

Literature: "Sarnak conjecture: What's new?" (Ferenczi - Kulaga Przymus - Lemanczyk)

### Motivation


#### Automatic sequence



$$n = 22 = (10110)_2,$$
  $u_{22} = 1$   
 $u = (u_2)_{22} = 01101001100101101001011001$ 

### Motivation

#### Automatic sequence




$$n = 22 = (10110)_2, u_{22} = 1$$

 $\mathbf{u} = (u_n)_{n \ge 0} = 01101001100101101001011001101001\dots$ 

### Motivation

#### Automatic sequence



$$n = 22 = (10110)_2, u_{22} = 1$$

$$\mathbf{u} = (u_n)_{n \ge 0} = 01101001100101101001011001101001\dots$$

Every automatic sequence  $(a_n)_{n\geq 0}$  fulfills the Sarnak Conjecture

Theorem 2 (M., 2016)


Under suitable (weak) conditions one also gets a Prime Number Theorem for automatic sequence.

Every automatic sequence  $(a_n)_{n\geq 0}$  fulfills the Sarnak Conjecture


#### Theorem 2 (M., 2016)

Under suitable (weak) conditions one also gets a Prime Number Theorem for automatic sequence.


For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.




For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.




For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.







For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.




For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.





For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.







For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.





For every strongly connected automaton A, there exists a naturally induced transducer  $\mathcal{T}_A$ . All other naturally induced transducers can be obtained by changing the order on the elements of Q.





### **Techniques**

## Use and adopt a framework of Mauduit and Rivat developed for the Rudin-Shapiro sequence.

- Carry Property: The contribution of high and low digits is "independent".
- Fourier Property: We say that U has the **Fourier property** if there exists  $\eta>0$  and c such that for all  $\lambda,\alpha$  and t

$$\left\|\frac{1}{k^{\lambda}}\sum_{m< k^{\lambda}}U(mk^{\alpha})e(mt)\right\|\leq ck^{-\eta\lambda}.$$

### **Techniques**

Use and adopt a framework of Mauduit and Rivat developed for the Rudin-Shapiro sequence.

- Carry Property: The contribution of high and low digits is "independent".
- Fourier Property: We say that U has the **Fourier property** if there exists  $\eta>0$  and c such that for all  $\lambda,\alpha$  and t

$$\left\|\frac{1}{k^{\lambda}}\sum_{m< k^{\lambda}}U(mk^{\alpha})e(mt)\right\|\leq ck^{-\eta\lambda}.$$

### **Techniques**

Use and adopt a framework of Mauduit and Rivat developed for the Rudin-Shapiro sequence.

- Carry Property: The contribution of high and low digits is "independent".
- Fourier Property: We say that U has the **Fourier property** if there exists  $\eta > 0$  and c such that for all  $\lambda, \alpha$  and t

$$\left\|\frac{1}{k^{\lambda}}\sum_{m< k^{\lambda}}U(mk^{\alpha})e(mt)\right\|\leq ck^{-\eta\lambda}.$$

## Zeckendorf Representation

#### Fibonacci numbers

$$F_0 = 0, F_1 = 1 \text{ and } F_{k+2} = F_{k+1} + F_k \text{ for } k \ge 0.$$

$$F_n = \frac{\varphi^n - (-\varphi)^{-n}}{\sqrt{5}},$$

where,  $\varphi$  is the golden ratio.

### Zeckendorf Representation

Every positive integer n admits a unique representation

$$n = \sum_{i>2} \varepsilon_i(n) F_i$$

where,  $\varepsilon_i(n) \in \{0,1\}$  and  $\varepsilon_i = 1 \Rightarrow \varepsilon_{i+1} = 0$ .

## Zeckendorf Representation

#### Fibonacci numbers

$$F_0 = 0, F_1 = 1$$
 and  $F_{k+2} = F_{k+1} + F_k$  for  $k \ge 0$ .

$$F_n = \frac{\varphi^n - (-\varphi)^{-n}}{\sqrt{5}},$$

where,  $\varphi$  is the golden ratio.

### Zeckendorf Representation

Every positive integer n admits a unique representation

$$n = \sum_{i \geq 2} \varepsilon_i(n) F_i$$

where,  $\varepsilon_i(n) \in \{0,1\}$  and  $\varepsilon_i = 1 \Rightarrow \varepsilon_{i+1} = 0$ .



## Zeckendorf Representation

#### Fibonacci numbers

$$F_0 = 0, F_1 = 1$$
 and  $F_{k+2} = F_{k+1} + F_k$  for  $k \ge 0$ .

$$F_n = \frac{\varphi^n - (-\varphi)^{-n}}{\sqrt{5}},$$

where,  $\varphi$  is the golden ratio.

#### Zeckendorf Representation

Every positive integer n admits a unique representation

$$n=\sum_{i\geq 2}\varepsilon_i(n)F_i,$$

where,  $\varepsilon_i(n) \in \{0,1\}$  and  $\varepsilon_i = 1 \Rightarrow \varepsilon_{i+1} = 0$ .

# Zeckendorf sum-of-digits Function

### Definition

We denote by

$$s_{\varphi}(n) = \sum_{i>2} \varepsilon_i(n)$$

the Zeckendorf sum-of-digits function.

We note that  $s_{\varphi}(n)$  is the least k such that n is the sum of k Fibonacci numbers.

# Zeckendorf sum-of-digits Function

#### Definition

We denote by

$$s_{\varphi}(n) = \sum_{i>2} \varepsilon_i(n)$$

the Zeckendorf sum-of-digits function.

We note that  $s_{\varphi}(n)$  is the least k such that n is the sum of k Fibonacci numbers.

## Main Result

### Theorem (Drmota, M., Spiegelhofer, 2017)

Let  $s_{\varphi}(n)$  be the Zeckendorf sum-of-digits function and m(n) a bounded multiplicative function. Then we have

$$\sum_{n\leq N} (-1)^{s_{\varphi}(n)} m(n) = o(N) \qquad (N\to\infty).$$

This implies that the Zeckendorf sum-of-digits function is orthogonal to the Möbius function.

## Main Result

### Theorem (Drmota, M., Spiegelhofer, 2017)

Let  $s_{\varphi}(n)$  be the Zeckendorf sum-of-digits function and m(n) a bounded multiplicative function. Then we have

$$\sum_{n\leq N} (-1)^{s_{\varphi}(n)} m(n) = o(N) \qquad (N\to\infty).$$

This implies that the Zeckendorf sum-of-digits function is orthogonal to the Möbius function.

### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if w is a fixed point of  $\sigma$ , i.e.  $\sigma(w) = w$ , then w is a k-automatic sequence.

### Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

### Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$\textit{E} = \{0,1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E=\{0,1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E=\{0,1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

## Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

01101001100101101001011001101001

#### Definition

Let E be a finite set and  $\sigma$  a k-uniform morphism such that  $\sigma(E) \subseteq E^k$ . Then if  $\mathbf{w}$  is a fixed point of  $\sigma$ , i.e.  $\sigma(\mathbf{w}) = \mathbf{w}$ , then  $\mathbf{w}$  is a k-automatic sequence.

### Example (Thue-Morse)

$$E = \{0, 1\}$$

$$\sigma(0) = 01$$

$$\sigma(1) = 10$$

### A Morphism

$$a \mapsto ab$$

$$b \mapsto c$$

$$c \mapsto cd$$

$$d \mapsto a$$
.

This gives the sequence  $(-1)^{s_{\varphi}(n)}$  under the coding  $\tau(a) = \tau(d) = 1, \tau(b) = \tau(c) = -1.$ 

This is one of the first examples of a substitution with non-constant length to be orthogonal to the Möbius function.

### A Morphism

$$a \mapsto ab$$

$$b \mapsto c$$

$$c \mapsto cd$$

$$d \mapsto a$$
.

This gives the sequence  $(-1)^{s_{\varphi}(n)}$  under the coding  $\tau(a) = \tau(d) = 1, \tau(b) = \tau(c) = -1.$ 

This is one of the first examples of a substitution with non-constant length to be orthogonal to the Möbius function.

### A Morphism

$$a \mapsto ab$$

$$b \mapsto c$$

$$c \mapsto cd$$

$$d \mapsto a$$
.


This gives the sequence  $(-1)^{s_{\varphi}(n)}$  under the coding  $\tau(a) = \tau(d) = 1, \tau(b) = \tau(c) = -1.$ 

This is one of the first examples of a substitution with non-constant length to be orthogonal to the Möbius function.

## A FAO

We use as input the Zeckendorf representation of n, i.e.

$$\varepsilon_k(n), \ldots, \varepsilon_0(n)$$
:



## Plan of the Proof

Use the Daboussi-Kátai Criterion to reduce the problem to

$$\sum_{n\leq N} (-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)} = o(N),$$

### for all different primes p, q.

• Use a generating function approach and "quasi-additivity" of  $(-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)}$  to reduce this to:

$$s_{\varphi}(pn_0) \not\equiv s_{\varphi}(qn_0) \bmod 2 \tag{1}$$

for some  $n_0$ 

• Show (1).



## Plan of the Proof

Use the Daboussi-Kátai Criterion to reduce the problem to

$$\sum_{n\leq N} (-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)} = o(N),$$

for all different primes p, q.

• Use a generating function approach and "quasi-additivity" of  $(-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)}$  to reduce this to:

$$s_{\varphi}(pn_0) \not\equiv s_{\varphi}(qn_0) \bmod 2 \tag{1}$$

for some  $n_0$ .

• Show (1).



## Plan of the Proof

Use the Daboussi-Kátai Criterion to reduce the problem to

$$\sum_{n\leq N} (-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)} = o(N),$$

for all different primes p, q.

• Use a generating function approach and "quasi-additivity" of  $(-1)^{s_{\varphi}(pn)+s_{\varphi}(qn)}$  to reduce this to:

$$s_{\varphi}(pn_0) \not\equiv s_{\varphi}(qn_0) \bmod 2 \tag{1}$$

for some  $n_0$ .

• Show (1).



Suppose that  $(x_n)$  is a bounded complex valued sequence with values in a finite set and that for every pair (p, q) of different prime numbers we have

$$\sum_{n\leq N} x_{pn} \overline{x_{qn}} = o(N).$$

Then for all bounded multiplicative functions  $\mathit{m}(\mathit{n})$  it follows that

$$\sum_{n \le N} x_n m(n) = o(N).$$

- Sum of digits in integer base: Dartyge Tennenbaum
- Rudin-Shapiro sequence: Tao, Mauduit-Rivat



Suppose that  $(x_n)$  is a bounded complex valued sequence with values in a finite set and that for every pair (p, q) of different prime numbers we have

$$\sum_{n\leq N} x_{pn} \overline{x_{qn}} = o(N).$$

Then for all bounded multiplicative functions m(n) it follows that

$$\sum_{n \le N} x_n m(n) = o(N).$$

- Sum of digits in integer base: Dartyge Tennenbaum
- Rudin-Shapiro sequence: Tao, Mauduit-Rivat



Suppose that  $(x_n)$  is a bounded complex valued sequence with values in a finite set and that for every pair (p, q) of different prime numbers we have

$$\sum_{n\leq N} x_{pn} \overline{x_{qn}} = o(N).$$

Then for all bounded multiplicative functions m(n) it follows that

$$\sum_{n\leq N}x_nm(n)=o(N).$$

- Sum of digits in integer base: Dartyge Tennenbaum
- Rudin-Shapiro seguence: Tao, Mauduit-Rivat



Suppose that  $(x_n)$  is a bounded complex valued sequence with values in a finite set and that for every pair (p, q) of different prime numbers we have

$$\sum_{n\leq N} x_{pn} \overline{x_{qn}} = o(N).$$

Then for all bounded multiplicative functions m(n) it follows that

$$\sum_{n\leq N}x_nm(n)=o(N).$$

- Sum of digits in integer base: Dartyge Tennenbaum
- Rudin-Shapiro sequence: Tao, Mauduit-Rivat



### Definition

We say that  $n_1$  and  $n_2$  are r-separated at position k if  $\varepsilon_i(n_1) = 0$  for  $i \ge k - r$  and  $\varepsilon_i(n_2) = 0$  for  $i \le k + r$ .

### Example:

$$n_1 = 4 \Rightarrow 0000101$$
  
 $n_2 = 29 \Rightarrow 1010000$ 

### Definition

We say that  $n_1$  and  $n_2$  are r-separated at position k if  $\varepsilon_i(n_1) = 0$  for  $i \ge k - r$  and  $\varepsilon_i(n_2) = 0$  for  $i \le k + r$ .

### Example:

$$n_1=4\Rightarrow 0000101$$

$$n_2 = 29 \Rightarrow 1010000$$

### Definition (for integer base by Kropf, Wagner)

We call a function f(n) quasi-additive (with respect to the Zeckendorf expansion) if there exists  $r \ge 0$  such that

•

$$f(n_1 + n_2) = f(n_1) + f(n_2)$$

for all integers  $n_1$ ,  $n_2$  that are r separated.

•  $f(n_1) = f(n_2)$  if the Zeckendorf expansion of  $n_1$  and  $n_2$  coincide up to "shifts".

### Definition (for integer base by Kropf, Wagner)

We call a function f(n) quasi-additive (with respect to the Zeckendorf expansion) if there exists  $r \ge 0$  such that

•

$$f(n_1 + n_2) = f(n_1) + f(n_2)$$

for all integers  $n_1$ ,  $n_2$  that are r separated.

•  $f(n_1) = f(n_2)$  if the Zeckendorf expansion of  $n_1$  and  $n_2$  coincide up to "shifts".

Let  $q > p \ge 2$  and  $f(n) = s_{\varphi}(pn) + s_{\varphi}(qn)$ . Then f(n) is quasi-additive with respect to the Zeckendorf expansion.

Proof (Sketch):

It suffices to work with  $s_{\varphi}(mn)$  as the sum of quasi-additive functions is again quasi-additive.

$$n_1 < F_{k-r} \Rightarrow mn_1 < F_k.$$

$$\varepsilon_i(n_2) = 0 \forall i < k + r \Rightarrow \varepsilon_i(mn_2) = 0 \forall i < k.$$

Let  $q > p \ge 2$  and  $f(n) = s_{\varphi}(pn) + s_{\varphi}(qn)$ . Then f(n) is quasi-additive with respect to the Zeckendorf expansion.

## Proof (Sketch):

It suffices to work with  $s_{\varphi}(mn)$  as the sum of quasi-additive functions is again quasi-additive.

Choose r such that  $\varphi^{r-1} < m$ .  $n_1 < F_{k-r} \Rightarrow mn_1 < F_k$ .  $\varepsilon_i(n_2) = 0 \forall i < k + r \Rightarrow \varepsilon_i(mn_2) = 0 \forall i < k$ .

Let  $q > p \ge 2$  and  $f(n) = s_{\varphi}(pn) + s_{\varphi}(qn)$ . Then f(n) is quasi-additive with respect to the Zeckendorf expansion.

Proof (Sketch):

It suffices to work with  $s_{\varphi}(mn)$  as the sum of quasi-additive functions is again quasi-additive.

$$n_1 < F_{k-r} \Rightarrow mn_1 < F_k.$$
  

$$\varepsilon_i(n_2) = 0 \forall i < k+r \Rightarrow \varepsilon_i(mn_2) = 0 \forall i < k.$$

Let  $q > p \ge 2$  and  $f(n) = s_{\varphi}(pn) + s_{\varphi}(qn)$ . Then f(n) is quasi-additive with respect to the Zeckendorf expansion.

Proof (Sketch):

It suffices to work with  $s_{\varphi}(mn)$  as the sum of quasi-additive functions is again quasi-additive.

$$n_1 < F_{k-r} \Rightarrow mn_1 < F_k.$$

$$\varepsilon_i(n_2) = 0 \forall i < k + r \Rightarrow \varepsilon_i(mn_2) = 0 \forall i < k.$$

Let  $q > p \ge 2$  and  $f(n) = s_{\varphi}(pn) + s_{\varphi}(qn)$ . Then f(n) is quasi-additive with respect to the Zeckendorf expansion.

Proof (Sketch):

It suffices to work with  $s_{\varphi}(mn)$  as the sum of quasi-additive functions is again quasi-additive.

$$\textit{n}_1 < \textit{F}_{k-r} \Rightarrow \textit{mn}_1 < \textit{F}_k.$$

$$\varepsilon_i(n_2) = 0 \forall i < k + r \Rightarrow \varepsilon_i(mn_2) = 0 \forall i < k.$$

## Generating Functions Approach

Let f be a quasi-additive function and

$$H(x,z) := \sum_{k \geq 3} x^k \sum_{F_{k-1} \leq n < F_k} z^{f(n)}.$$

Note that

$$[x^k]H(x,-1) = \sum_{F_{k-1} \le n < F_k} (-1)^{s_{\varphi}(pn) + s_{\varphi}(qn)}$$

Let  $\mathcal B$  be the set of integers n whose Zeckendorf expansion ends with exactly r zeros and that can not be decomposed into positive, r-separated summands. Let

$$B(x,z) = \sum_{n \in \mathcal{B}} x^{\ell(n)} z^{f(n)},$$

where  $\ell(n) = k$  if  $F_{k-1} \le n < F_k$ .



## Generating Functions Approach

Let f be a quasi-additive function and

$$H(x,z) := \sum_{k \geq 3} x^k \sum_{F_{k-1} \leq n < F_k} z^{f(n)}.$$

Note that

$$[x^k]H(x,-1) = \sum_{F_{k-1} \le n < F_k} (-1)^{s_{\varphi}(pn) + s_{\varphi}(qn)}.$$

Let  $\mathcal B$  be the set of integers n whose Zeckendorf expansion ends with exactly r zeros and that can not be decomposed into positive, r-separated summands. Let

$$B(x,z) = \sum_{n \in \mathcal{B}} x^{\ell(n)} z^{f(n)},$$

where  $\ell(n) = k$  if  $F_{k-1} \le n < F_k$ .



## Generating Functions Approach

Let f be a quasi-additive function and

$$H(x,z) := \sum_{k \geq 3} x^k \sum_{F_{k-1} \leq n < F_k} z^{f(n)}.$$

Note that

$$[x^k]H(x,-1) = \sum_{F_{k-1} \le n < F_k} (-1)^{s_{\varphi}(pn) + s_{\varphi}(qn)}.$$

Let  $\mathcal B$  be the set of integers n whose Zeckendorf expansion ends with exactly r zeros and that can not be decomposed into positive, r-separated summands. Let

$$B(x,z) = \sum_{n \in \mathcal{B}} x^{\ell(n)} z^{f(n)},$$

where  $\ell(n) = k$  if  $F_{k-1} \le n < F_k$ .



$$H(x,z) = \frac{1}{1-x} \frac{1}{1-B(x,z) \frac{x^{2r+1}}{1-x}} B'(x,z)$$
$$= \frac{B'(x,z)}{1-x-x^{2r+1}B(x,z)}.$$

The dominant singularity of H(x,1) is at  $x_0 = \frac{1}{\varphi}$ . This is due to the fact that  $x = x_0$  is a solution fo

$$x + x^{2r+1}B(x,1) = 1$$

$$x + x^{2r+1}B(x, -1) = 1.$$



$$H(x,z) = \frac{1}{1-x} \frac{1}{1-B(x,z) \frac{x^{2r+1}}{1-x}} B'(x,z)$$
$$= \frac{B'(x,z)}{1-x-x^{2r+1}B(x,z)}.$$

The dominant singularity of H(x,1) is at  $x_0 = \frac{1}{\varphi}$ .

This is due to the fact that  $x = x_0$  is a solution for

$$x + x^{2r+1}B(x,1) = 1$$

$$x + x^{2r+1}B(x, -1) = 1.$$



$$H(x,z) = \frac{1}{1-x} \frac{1}{1-B(x,z) \frac{x^{2r+1}}{1-x}} B'(x,z)$$
$$= \frac{B'(x,z)}{1-x-x^{2r+1}B(x,z)}.$$

The dominant singularity of H(x,1) is at  $x_0 = \frac{1}{\varphi}$ . This is due to the fact that  $x = x_0$  is a solution for

$$x + x^{2r+1}B(x,1) = 1.$$

$$x + x^{2r+1}B(x, -1) = 1.$$



$$H(x,z) = \frac{1}{1-x} \frac{1}{1-B(x,z) \frac{x^{2r+1}}{1-x}} B'(x,z)$$
$$= \frac{B'(x,z)}{1-x-x^{2r+1}B(x,z)}.$$

The dominant singularity of H(x,1) is at  $x_0 = \frac{1}{\varphi}$ . This is due to the fact that  $x = x_0$  is a solution for

$$x + x^{2r+1}B(x,1) = 1.$$

$$x + x^{2r+1}B(x, -1) = 1.$$

It remains to find *n* such that

$$s_{\varphi}(pn) + s_{\varphi}(qn) \equiv 1 \mod 2.$$

The key point is to find  $n_1$ ,  $n_2$  such that

$$egin{aligned} s_arphi(pn_1)+s_arphi(pn_2)&\equiv s_arphi(p(n_1+n_2))\ ext{mod}\ 2\ s_arphi(qn_1)+s_arphi(qn_2)&\equiv s_arphi(q(n_1+n_2))+1\ ext{mod}\ 2. \end{aligned}$$

It remains to find *n* such that

$$s_{\varphi}(pn) + s_{\varphi}(qn) \equiv 1 \mod 2.$$

The key point is to find  $n_1$ ,  $n_2$  such that

$$\begin{split} s_{\varphi}(pn_1) + s_{\varphi}(pn_2) &\equiv s_{\varphi}(p(n_1+n_2)) \bmod 2 \\ s_{\varphi}(qn_1) + s_{\varphi}(qn_2) &\equiv s_{\varphi}(q(n_1+n_2)) + 1 \bmod 2. \end{split}$$