

Normal Subsequences of Automatic Sequences

Clemens Müllner


8. Jun 2017

Automatic Sequences

Definition (Automaton - DFA)

$$A = (Q, \Sigma = \{0, \dots, k-1\}, \delta, q_0, \tau)$$

Example (Thue-Morse sequence)



$$n = 22 = (10110)_2, \quad u_{22} = 1$$


$$\mathbf{u} = (u_n)_{n \geq 0} = 01101001100101101001011001101001 \dots$$

Automatic Sequences

Definition (Automaton - DFA)

$$A = (Q, \Sigma = \{0, \dots, k-1\}, \delta, q_0, \tau)$$

Example (Thue-Morse sequence)



$$n = 22 = (10110)_2, \quad u_{22} = 1$$


$$\mathbf{u} = (u_n)_{n \geq 0} = 01101001100101101001011001101001 \dots$$

Automatic Sequences

Definition (Automaton - DFA)

$$A = (Q, \Sigma = \{0, \dots, k-1\}, \delta, q_0, \tau)$$

Example (Thue-Morse sequence)



$$n = 22 = (10110)_2, \quad u_{22} = 1$$


$$\mathbf{u} = (u_n)_{n \geq 0} = 01101001100101101001011001101001 \dots$$

Automatic Sequences

Definition (Automaton - DFA)

$$A = (Q, \Sigma = \{0, \dots, k-1\}, \delta, q_0, \tau)$$

Example (Thue-Morse sequence)



$$n = 22 = (10110)_2, \quad u_{22} = 1$$

$$\mathbf{u} = (u_n)_{n \geq 0} = 01101001100101101001011001101001 \dots$$

Examples of Automatic Sequences

- Periodic sequences.
- q -additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n)) \text{ and } f(0) = 0.$$

- q -block-additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n), \dots, \varepsilon_{j+r}(n)) \text{ and } f(0, \dots, 0) = 0.$$

Examples of Automatic Sequences

- Periodic sequences.
- q -additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n)) \text{ and } f(0) = 0.$$

- q -block-additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n), \dots, \varepsilon_{j+r}(n)) \text{ and } f(0, \dots, 0) = 0.$$

Examples of Automatic Sequences

- Periodic sequences.
- q -additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n)) \text{ and } f(0) = 0.$$

- q -block-additive function modulo m : $u_n = f(n) \bmod m$

$$f(n) = \sum_{j \geq 0} f(\varepsilon_j(n), \dots, \varepsilon_{j+r}(n)) \text{ and } f(0, \dots, 0) = 0.$$

Properties of Automatic Sequences

- For every automatic sequence \mathbf{u} there exists the logarithmic density

$$\text{logdens}(\mathbf{u}, a) = \lim_{N \rightarrow \infty} \frac{1}{\log(N)} \sum_{1 \leq n \leq N} \frac{1}{n} \mathbf{1}_{[u_n=a]}.$$

- The subword complexity p_k of an automatic sequence is (at most) linear.
- Every subsequence $(u_{an+b})_{n \geq 0}$ along an arithmetic progression of an automatic sequence $(u_n)_{n \geq 0}$ is again automatic.
- Let $u^{(1)}(n), \dots, u^{(j)}(n)$ be automatic sequences. Then $u(n) = f(u^{(1)}(n), \dots, u^{(j)}(n))$ is again automatic.

Properties of Automatic Sequences

- For every automatic sequence \mathbf{u} there exists the logarithmic density

$$\text{logdens}(\mathbf{u}, a) = \lim_{N \rightarrow \infty} \frac{1}{\log(N)} \sum_{1 \leq n \leq N} \frac{1}{n} \mathbf{1}_{[u_n=a]}.$$

- The subword complexity p_k of an automatic sequence is (at most) linear.
- Every subsequence $(u_{an+b})_{n \geq 0}$ along an arithmetic progression of an automatic sequence $(u_n)_{n \geq 0}$ is again automatic.
- Let $u^{(1)}(n), \dots, u^{(j)}(n)$ be automatic sequences. Then $u(n) = f(u^{(1)}(n), \dots, u^{(j)}(n))$ is again automatic.

Properties of Automatic Sequences

- For every automatic sequence \mathbf{u} there exists the logarithmic density

$$\text{logdens}(\mathbf{u}, a) = \lim_{N \rightarrow \infty} \frac{1}{\log(N)} \sum_{1 \leq n \leq N} \frac{1}{n} \mathbf{1}_{[u_n=a]}.$$

- The subword complexity p_k of an automatic sequence is (at most) linear.
- Every subsequence $(u_{an+b})_{n \geq 0}$ along an arithmetic progression of an automatic sequence $(u_n)_{n \geq 0}$ is again automatic.
- Let $u^{(1)}(n), \dots, u^{(j)}(n)$ be automatic sequences. Then $u(n) = f(u^{(1)}(n), \dots, u^{(j)}(n))$ is again automatic.

Properties of Automatic Sequences

- For every automatic sequence \mathbf{u} there exists the logarithmic density

$$\text{logdens}(\mathbf{u}, a) = \lim_{N \rightarrow \infty} \frac{1}{\log(N)} \sum_{1 \leq n \leq N} \frac{1}{n} \mathbf{1}_{[u_n=a]}.$$

- The subword complexity p_k of an automatic sequence is (at most) linear.
- Every subsequence $(u_{an+b})_{n \geq 0}$ along an arithmetic progression of an automatic sequence $(u_n)_{n \geq 0}$ is again automatic.
- Let $u^{(1)}(n), \dots, u^{(j)}(n)$ be automatic sequences. Then $u(n) = f(u^{(1)}(n), \dots, u^{(j)}(n))$ is again automatic.

General Idea

- Start with an automatic sequence u_n that is uniformly distributed on the output alphabet.
- Consider a relatively sparse subsequence u_{n_k} that has the same asymptotic frequencies. (The size of the gaps needs to increase sufficiently fast.)
- This subsequence should be pseudo-random (or normal) on the output alphabet.

General Idea

- Start with an automatic sequence u_n that is uniformly distributed on the output alphabet.
- Consider a relatively sparse subsequence u_{n_k} that has the same asymptotic frequencies. (The size of the gaps needs to increase sufficiently fast.)
- This subsequence should be pseudo-random (or normal) on the output alphabet.

General Idea

- Start with an automatic sequence u_n that is uniformly distributed on the output alphabet.
- Consider a relatively sparse subsequence u_{n_k} that has the same asymptotic frequencies. (The size of the gaps needs to increase sufficiently fast.)
- This subsequence should be pseudo-random (or normal) on the output alphabet.

Thue-Morse sequence along Piatetski-Shapiro sequence $\lfloor n^c \rfloor$

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (1995, 2005), Spiegelhofer(2014,2015+)

$1 < c < 3/2$:

$$\#\{0 \leq n < N : t_{\lfloor n^c \rfloor} = 0\} \approx \frac{N}{2}.$$

Thue-Morse sequence along Piatetski-Shapiro sequence $\lfloor n^c \rfloor$

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (1995, 2005), Spiegelhofer(2014,2015+)

$1 < c < 3/2$:

$$\#\{0 \leq n < N : t_{\lfloor n^c \rfloor} = 0\} \approx \frac{N}{2}.$$

Thue-Morse sequence along Piatetski-Shapiro sequence $\lfloor n^c \rfloor$

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (1995, 2005), Spiegelhofer(2014,2015+)

$1 < c < 3/2$:

$$\#\{0 \leq n < N : t_{\lfloor n^c \rfloor} = 0\} \approx \frac{N}{2}.$$

Subsequences along $\lfloor n^c \rfloor$

Theorem (Deshouillers, Drmota and Morgenbesser, 2012)

Let u_n be a k -automatic sequence (on an alphabet \mathcal{A}) and

$$1 < c < 7/5.$$

Then for each $a \in \mathcal{A}$ the asymptotic density $\text{dens}(u_{\lfloor n^c \rfloor}, a)$ of a in the subsequence u_{n^c} exists if and only if the asymptotic density of a in u_n exists and we have

$$\text{dens}(u_{\lfloor n^c \rfloor}, a) = \text{dens}(u_n, a).$$

Thue-Morse sequence along squares

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2009):

$$\#\{0 \leq n < N : t_{n^2} = 0\} \approx \frac{N}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along squares

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2009):

$$\#\{0 \leq n < N : t_{n^2} = 0\} \approx \frac{N}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along squares

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2009):

$$\#\{0 \leq n < N : t_{n^2} = 0\} \approx \frac{N}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along squares

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2009):

$$\#\{0 \leq n < N : t_{n^2} = 0\} \approx \frac{N}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Subsequences along squares

Theorem (M., 2017+)

Let u_n be a k -automatic sequence (on an alphabet \mathcal{A}) generated by a strongly connected automaton such that a initial state is mapped to itself under 0. Then for each $a \in \mathcal{A}$ the asymptotic density

$$\text{dens}(u_{n^2}, a)$$

exists (and can be computed).

Thue-Morse sequence along primes

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2010):

$$\#\{0 \leq p < N : t_p = 0\} \approx \frac{\pi(N)}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along primes

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2010):

$$\#\{0 \leq p < N : t_p = 0\} \approx \frac{\pi(N)}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along primes

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2010):

$$\#\{0 \leq p < N : t_p = 0\} \approx \frac{\pi(N)}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Thue-Morse sequence along primes

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2010):

$$\#\{0 \leq p < N : t_p = 0\} \approx \frac{\pi(N)}{2}.$$

Solution of a Conjecture of Gelfond (1968).

Subsequences along primes

Theorem (M., 2017+)

Let u_n be a k -automatic sequence (on an alphabet \mathcal{A}) generated by a strongly connected automaton such that a initial state is mapped to itself under 0. Then for each $a \in \mathcal{A}$ the asymptotic density

$$\text{dens}(u_{p_n}, a)$$

exists, where p_n denotes the n -th prime number (and can be computed).

Sarnak Conjecture for automatic sequences

Theorem (M., 2016)

Let u_n be a complex-valued automatic sequence.

Then we have

$$\sum_{n \leq N} u_n \mu(n) = o(N),$$

where $\mu(n)$ denotes the Möbius function.

This generalizes several results by Dartyge and Tenenbaum (Thue-Morse); Mauduit and Rivat (Rudin-Shapiro); Tao (Rudin-Shapiro); Drmota (invertible); Ferenczi, Kulaga-Przymus, Lemanczyk, and Mauduit (invertible); Deshoulliers, Drmota and M. (synchronizing).

Thue-Morse sequence along squares

$p_k^{(2)}$... subword complexity of $(t_{n^2})_{n \geq 0}$.

Conjecture (Allouche and Shallit, 2003)

$$p_k^{(2)} = 2^k$$

Equivalently: every block $B \in \{0, 1\}^k$, $k \geq 1$, appears in $(t_{n^2})_{n \geq 0}$.

(Moshe, 2007): $p_k^{(2)} = 2^k$.

But what can be said about the frequency of a given block?

Thue-Morse sequence along squares

$p_k^{(2)}$... subword complexity of $(t_{n^2})_{n \geq 0}$.

Conjecture (Allouche and Shallit, 2003)

$$p_k^{(2)} = 2^k$$

Equivalently: every block $B \in \{0, 1\}^k$, $k \geq 1$, appears in $(t_{n^2})_{n \geq 0}$.

(Moshe, 2007): $p_k^{(2)} = 2^k$.

But what can be said about the frequency of a given block?

Thue-Morse sequence along squares

$p_k^{(2)}$... subword complexity of $(t_{n^2})_{n \geq 0}$.

Conjecture (Allouche and Shallit, 2003)

$$p_k^{(2)} = 2^k$$

Equivalently: every block $B \in \{0, 1\}^k$, $k \geq 1$, appears in $(t_{n^2})_{n \geq 0}$.

(Moshe, 2007): $p_k^{(2)} = 2^k$.

But what can be said about the frequency of a given block?

Thue-Morse sequence along squares

$p_k^{(2)}$... subword complexity of $(t_{n^2})_{n \geq 0}$.

Conjecture (Allouche and Shallit, 2003)

$$p_k^{(2)} = 2^k$$

Equivalently: every block $B \in \{0, 1\}^k$, $k \geq 1$, appears in $(t_{n^2})_{n \geq 0}$.

(Moshe, 2007): $p_k^{(2)} = 2^k$.

But what can be said about the frequency of a given block?

Definition

A sequence $(u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}$ is *normal* if for any $k \in \mathbb{N}$ and any $B = (b_0, \dots, b_{k-1}) \in \{0, 1\}^k$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{i < N, u_i = b_0, \dots, u_{i+k-1} = b_{k-1}\} = \frac{1}{2^k}.$$

There are only few (known) explicit examples of normal sequences.

Definition

A sequence $(u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}$ is *normal* if for any $k \in \mathbb{N}$ and any $B = (b_0, \dots, b_{k-1}) \in \{0, 1\}^k$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{i < N, u_i = b_0, \dots, u_{i+k-1} = b_{k-1}\} = \frac{1}{2^k}.$$

There are only few (known) explicit examples of normal sequences.

Normal subsequences of the Thue-Morse sequence

Theorem (Drmota + Mauduit + Rivat, 2013+)

The sequence (t_{n^2}) is normal.

Theorem (M. + Spiegelhofer, 2015+)

Suppose that $1 < c < 3/2$. Then the sequence $(t_{\lfloor n^c \rfloor})$ is normal.

Normal subsequences of the Thue-Morse sequence

Theorem (Drmota + Mauduit + Rivat, 2013+)

The sequence (t_{n^2}) is normal.

Theorem (M. + Spiegelhofer, 2015+)

Suppose that $1 < c < 3/2$. Then the sequence $(t_{\lfloor n^c \rfloor})$ is normal.

Normal subsequences

Theorem (M., 2017+)

Let $f(n)$ be a block-additive function and $u_n = f(n) \bmod m$ an automatic sequence which is uniformly distributed on the alphabet $\{0, \dots, m-1\}$ along arithmetic subsequences.

Then the sequence $(u_{\lfloor n^c \rfloor})_{n \geq 0}$ is normal for all c with $1 < c < 4/3$. Furthermore, $(u_{n^2})_{n \geq 0}$ is normal.

Conjecture (1)

Suppose that $c > 1$ and $c \notin \mathbb{Z}$. Then for every automatic sequence u_n (on an alphabet \mathcal{A}) the asymptotic density $\text{dens}(u_{\lfloor n^c \rfloor}, a)$ of $a \in \mathcal{A}$ in the subsequence $(u_{\lfloor n^c \rfloor})$ exists if and only if the asymptotic density of a in u_n exists and we have up to periodic behavior

$$\begin{aligned} \lim_{N \rightarrow \infty} \#\{n < N, u_{\lfloor n^c \rfloor} = b_0, \dots, u_{\lfloor (n+k-1)^c \rfloor} = b_{k-1}\} \\ = \text{dens}(u_n, b_0) \cdots \text{dens}(u_n, b_{k-1}) \end{aligned}$$

for every $k \geq 1$ and for all $b_0, \dots, b_{k-1} \in \mathcal{A}$.

Conjecture (2)

Let $P(x)$ be a positive integer valued polynomial and u_n an automatic sequence generated by a strongly connected automaton. Then, for every $a \in \mathcal{A}$ the densities $\delta_a = \text{dens}(u_{P(n)}, a)$ exists and we have (up to periodic behavior)

$$\begin{aligned} \lim_{N \rightarrow \infty} \#\{n < N, u_{P(n)} = b_0, \dots, u_{P(n+k-1)} = b_{k-1}\} \\ = \delta_{b_0} \cdots \delta_{b_{k-1}} \end{aligned}$$

for every $k \geq 1$ and for all $b_0, \dots, b_{k-1} \in \mathcal{A}$.

Let u_n be an automatic sequence and $\phi(n)$ a positive sequence such that $\phi(n)/n$ is non-decreasing.

What can be said about $u_{\lfloor \phi(n) \rfloor}$?

- We cannot expect general results for exponentially growing sequences $\phi(n)$.
- If $\phi(n) = an + b$ with integers a, b . Then $u_{\phi(n)}$ is again an automatic sequence.
- If $\phi(n) = n \log_2(n)$ then $t_{\lfloor \phi(n) \rfloor}$ behaves like the Thue-Morse sequence t_n , but the density for blocks of length 2 does not exist. (Deshouillers + Drmota + Morgenbesser (2012))

Let u_n be an automatic sequence and $\phi(n)$ a positive sequence such that $\phi(n)/n$ is non-decreasing.

What can be said about $u_{\lfloor \phi(n) \rfloor}$?

- We cannot expect general results for exponentially growing sequences $\phi(n)$.
- If $\phi(n) = an + b$ with integers a, b . Then $u_{\phi(n)}$ is again an automatic sequence.
- If $\phi(n) = n \log_2(n)$ then $t_{\lfloor \phi(n) \rfloor}$ behaves like the Thue-Morse sequence t_n , but the density for blocks of length 2 does not exist. (Deshouillers + Drmota + Morgenbesser (2012))

Let u_n be an automatic sequence and $\phi(n)$ a positive sequence such that $\phi(n)/n$ is non-decreasing.

What can be said about $u_{\lfloor \phi(n) \rfloor}$?

- We cannot expect general results for exponentially growing sequences $\phi(n)$.
- If $\phi(n) = an + b$ with integers a, b . Then $u_{\phi(n)}$ is again an automatic sequence.
- If $\phi(n) = n \log_2(n)$ then $t_{\lfloor \phi(n) \rfloor}$ behaves like the Thue-Morse sequence t_n , but the density for blocks of length 2 does not exist. (Deshouillers + Drmota + Morgenbesser (2012))

Let u_n be an automatic sequence and $\phi(n)$ a positive sequence such that $\phi(n)/n$ is non-decreasing.

What can be said about $u_{\lfloor \phi(n) \rfloor}$?

- We cannot expect general results for exponentially growing sequences $\phi(n)$.
- If $\phi(n) = an + b$ with integers a, b . Then $u_{\phi(n)}$ is again an automatic sequence.
- If $\phi(n) = n \log_2(n)$ then $t_{\lfloor \phi(n) \rfloor}$ behaves like the Thue-Morse sequence t_n , but the density for blocks of length 2 does not exist. (Deshouillers + Drmota + Morgenbesser (2012))

- Rewrite the statement in terms of exponential sums.
E.g. $\text{dens}(t_{n^2}, 0) = 1/2$ holds if

$$\left| \sum_{n \leq N} e\left(\frac{s_2(n^2)}{2}\right) \right| = o(N),$$

where $e(x) = \exp(2\pi ix)$.

- Use independence of „high“ and „low“ digits.
- Statement involving the discrete Fourier transform

$$F_\lambda(h, \alpha) = \frac{1}{2^\lambda} \sum_{u < 2^\lambda} e(\alpha s_2(u) - hu2^{-\lambda}).$$

- Continuation of the example:

$$|F_\lambda(h, 1/2)| \leq 2^{-\eta m} |F_{\lambda-m}(h, 1/2)|,$$

for some m and $\eta > 0$.

- Rewrite the statement in terms of exponential sums.
E.g. $\text{dens}(t_{n^2}, 0) = 1/2$ holds if

$$\left| \sum_{n \leq N} e\left(\frac{s_2(n^2)}{2}\right) \right| = o(N),$$

where $e(x) = \exp(2\pi ix)$.

- Use independence of „high“ and „low“ digits.
- Statement involving the discrete Fourier transform

$$F_\lambda(h, \alpha) = \frac{1}{2^\lambda} \sum_{u < 2^\lambda} e(\alpha s_2(u) - hu2^{-\lambda}).$$

- Continuation of the example:

$$|F_\lambda(h, 1/2)| \leq 2^{-\eta m} |F_{\lambda-m}(h, 1/2)|,$$

for some m and $\eta > 0$.

- Rewrite the statement in terms of exponential sums.
E.g. $\text{dens}(t_{n^2}, 0) = 1/2$ holds if

$$\left| \sum_{n \leq N} e\left(\frac{s_2(n^2)}{2}\right) \right| = o(N),$$

where $e(x) = \exp(2\pi ix)$.

- Use independence of „high“ and „low“ digits.
- Statement involving the discrete Fourier transform

$$F_\lambda(h, \alpha) = \frac{1}{2^\lambda} \sum_{u < 2^\lambda} e(\alpha s_2(u) - hu2^{-\lambda}).$$

- Continuation of the example:

$$|F_\lambda(h, 1/2)| \leq 2^{-\eta m} |F_{\lambda-m}(h, 1/2)|,$$

for some m and $\eta > 0$.

- Rewrite the statement in terms of exponential sums.
E.g. $\text{dens}(t_{n^2}, 0) = 1/2$ holds if

$$\left| \sum_{n \leq N} e\left(\frac{s_2(n^2)}{2}\right) \right| = o(N),$$

where $e(x) = \exp(2\pi ix)$.

- Use independence of „high“ and „low“ digits.
- Statement involving the discrete Fourier transform

$$F_\lambda(h, \alpha) = \frac{1}{2^\lambda} \sum_{u < 2^\lambda} e(\alpha s_2(u) - hu2^{-\lambda}).$$



- Continuation of the example:

$$|F_\lambda(h, 1/2)| \leq 2^{-\eta m} |F_{\lambda-m}(h, 1/2)|,$$

for some m and $\eta > 0$.



Representation of automatic sequences

Example (Rudin-Shapiro)





Representation of automatic sequences

Example (Rudin-Shapiro)



Representation of automatic sequences


Example (Rudin-Shapiro)



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .


Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .



Example:



Theorem (M., 2016)

For every strongly connected automaton A , there exists a naturally induced transducer \mathcal{T}_A . All other naturally induced transducers can be obtained by changing the order on the elements of Q .

Example:



Fibonacci Base

Theorem (Drmotá, M., Spiegelhofer, 2017+)

Let $s_\varphi(n)$ be the Zeckendorf sum-of-digits function and $m(n)$ a bounded multiplicative function. Then we have

$$\sum_{n < N} (-1)^{s_\varphi(n)} m(n) = o(N) \quad (N \rightarrow \infty).$$

This implies that the Zeckendorf sum-of-digits function is orthogonal to the Möbius function.

Fibonacci Base

Theorem (Drmotá, M., Spiegelhofer, 2017+)

Let $s_\varphi(n)$ be the Zeckendorf sum-of-digits function and $m(n)$ a bounded multiplicative function. Then we have

$$\sum_{n < N} (-1)^{s_\varphi(n)} m(n) = o(N) \quad (N \rightarrow \infty).$$

This implies that the Zeckendorf sum-of-digits function is orthogonal to the Möbius function.