

AFFINE FRACTIONAL SOBOLEV AND ISOPERIMETRIC INEQUALITIES

JULIÁN HADDAD AND MONIKA LUDWIG

ABSTRACT. Sharp affine fractional Sobolev inequalities for functions on \mathbb{R}^n are established. For each $0 < s < 1$, the new inequalities are significantly stronger than (and directly imply) the sharp fractional Sobolev inequalities of Almgren and Lieb. In the limit as $s \rightarrow 1^-$, the new inequalities imply the sharp affine Sobolev inequality of Gaoyong Zhang. As a consequence, fractional Petty projection inequalities are obtained that are stronger than the fractional Euclidean isoperimetric inequalities and a natural conjecture for radial mean bodies is proved.

2020 AMS subject classification: 46E35 (35R11, 52A40)

1. INTRODUCTION

Almgren and Lieb [1] established Pólya–Szegő inequalities for fractional Sobolev norms. The equality case was settled by Frank and Seiringer [11]. A consequence is the fractional Sobolev inequality: for $0 < s < 1$, there is $\alpha_{n,s} > 0$ such that for $f \in W^{s,1}(\mathbb{R}^n)$,

$$(1) \quad \|f\|_{\frac{n}{n-s}} \leq \alpha_{n,s} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy,$$

and there is equality if and only if f is a constant multiple of the indicator function of a ball. Here $\|f\|_p$ denotes the L^p norm of f and $|\cdot|$ the Euclidean norm in \mathbb{R}^n while $W^{s,1}(\mathbb{R}^n)$ denotes the fractional Sobolev space of L^1 functions with finite right side in (1). By a result of Bourgain, Brezis, and Mironescu [3], for $f \in W^{1,1}(\mathbb{R}^n)$,

$$(2) \quad \lim_{s \rightarrow 1^-} \alpha_{n,s} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy = \frac{1}{n\omega_n^{1/n}} \int_{\mathbb{R}^n} |\nabla f(x)| dx,$$

where ω_n is the n -dimensional volume of the unit ball in \mathbb{R}^n and $W^{1,1}(\mathbb{R}^n)$ denotes the Sobolev space of L^1 functions f with weak L^1 gradient ∇f . Dávila [7] proved a result corresponding to (2) for $f \in BV(\mathbb{R}^n)$, the space of L^1 functions with bounded variation. Hence, the sharp L^1 Sobolev inequalities on $W^{1,1}(\mathbb{R}^n)$ and $BV(\mathbb{R}^n)$ follow from the fractional Sobolev inequalities (1).

Gaoyong Zhang [38] established a sharp affine Sobolev inequality that is significantly stronger than the classical L^1 Sobolev inequality. For $n \geq 2$ and C^1 functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ with compact support, Zhang [38] proved that

$$(3) \quad \|f\|_{\frac{n}{n-1}} \leq \frac{\omega_n}{2\omega_{n-1}} \left(\frac{1}{n} \int_{\mathbb{S}^{n-1}} \|\langle \nabla f(\cdot), \xi \rangle\|_1^{-n} d\xi \right)^{-1/n} \leq \frac{1}{n\omega_n^{1/n}} \int_{\mathbb{R}^n} |\nabla f(x)| dx.$$

Here $\langle \cdot, \cdot \rangle$ denotes the inner product and integration on the $(n-1)$ -dimensional unit sphere \mathbb{S}^{n-1} is with respect to the $(n-1)$ -dimensional Hausdorff measure. The

first inequality in (3) is now called the affine Sobolev inequality or the Sobolev–Zhang inequality. It is affine since both terms remain invariant under volume-preserving affine transformations. Tuo Wang [35] extended the Sobolev–Zhang inequality to L^1 functions of bounded variation and showed that there is equality in this generalized Sobolev–Zhang inequality precisely for constant multiples of indicator functions of ellipsoids.

The main aim of this paper is to establish affine fractional Sobolev inequalities for $0 < s < 1$ that are stronger than the fractional Sobolev inequalities in (1) and imply, in the limit as $s \rightarrow 1^-$, the Sobolev–Zhang inequality.

Theorem 1. *For $0 < s < 1$ and $f \in W^{s,1}(\mathbb{R}^n)$,*

$$\begin{aligned} \|f\|_{\frac{n}{n-s}} &\leq \alpha_{n,s} n \omega_n^{(n+s)/n} \left(\frac{1}{n} \int_{\mathbb{S}^{n-1}} \left(\int_0^\infty t^{-s} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 dt \right)^{-n/s} d\xi \right)^{-s/n} \\ &\leq \alpha_{n,s} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy. \end{aligned}$$

There is equality in the first inequality if and only if f is a constant multiple of the indicator function of an ellipsoid. There is equality in the second inequality if f is radially symmetric.

In order to prove Theorem 1 we introduce the s -fractional polar projection body $\Pi^{*,s}f$ associated to f , defined as the star body whose gauge function for $\xi \in \mathbb{S}^{n-1}$ is

$$(4) \quad \|\xi\|_{\Pi^{*,s}f}^s = \int_0^\infty t^{-s} \int_{\mathbb{R}^n} \left| \frac{f(x + t\xi) - f(x)}{t} \right| dx dt$$

(see Section 3 for details). The affine fractional Sobolev inequality now can be written as

$$(5) \quad \|f\|_{\frac{n}{n-s}} \leq \alpha_{n,s} n \omega_n^{(n+s)/n} |\Pi^{*,s}f|^{-s/n},$$

where $|\cdot|$ denotes n -dimensional Lebesgue measure. Since both sides of (5) are invariant under translations of f , and for volume-preserving linear transformations $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$,

$$\Pi^{*,s}(f \circ \phi^{-1}) = \phi \Pi^{*,s}f,$$

it follows that (5) is an affine inequality.

To show that Theorem 1 implies the Sobolev–Zhang inequality, we establish a result corresponding to (2). For $f \in W^{1,1}(\mathbb{R}^n)$, we will prove that

$$(6) \quad \lim_{s \rightarrow 1^-} (1-s) |\Pi^{*,s}f|^{-s/n} = 2 |\Pi^*f|^{-1/n},$$

where Π^*f is the polar projection body of f which is defined (with different notation) in [38] (also see [24]) for $\xi \in \mathbb{S}^{n-1}$ by

$$\|\xi\|_{\Pi^*f} = \frac{1}{2} \int_{\mathbb{R}^n} |\langle \nabla f(x), \xi \rangle| dx.$$

Moreover, we establish the corresponding statement for $f \in BV(\mathbb{R}^n)$ and also recover Wang’s generalized Sobolev–Zhang inequality (without characterization of the equality case) from Theorem 1 (see Section 4).

We prove Theorem 1 by first establishing affine fractional Pólya–Szegő inequalities as a consequence of the Riesz rearrangement inequality. For the equality case, we use anisotropic fractional Sobolev norms (introduced in [21]) and the equality case of the Riesz rearrangement inequality due to Burchard [5]. For $0 < s < 1$ and $E \subset \mathbb{R}^n$ of finite s -perimeter (see Section 2), we define the fractional polar projection body as $\Pi^{*,s}E = \Pi^{*,s}1_E$, where 1_E is the indicator function of E , and obtain from the affine fractional Pólya–Szegő inequalities that

$$(7) \quad \left(\frac{|\Pi^{*,s}E|}{|\Pi^{*,s}B^n|} \right)^{-s/n} \geq \left(\frac{|E|}{|B^n|} \right)^{(n-s)/n}$$

for $0 < s < 1$ and $E \subset \mathbb{R}^n$ of finite s -perimeter, where B^n is the Euclidean unit ball in \mathbb{R}^n . From these fractional Petty projection inequalities, Theorem 1 follows by a co-area formula for the fractional Sobolev norm. In the limit as $s \rightarrow 1^-$, we obtain from (7) the generalized Petty projection inequality by Zhang [38] and Wang [35]. In addition, we prove versions of our functional and geometric inequalities for Steiner symmetrization.

In Section 9, we connect fractional polar projection bodies to radial mean bodies (introduced by Gardner and Zhang [13]) and obtain fractional Zhang inequalities. Moreover, we establish new affine isoperimetric inequalities for radial mean bodies, thereby proving a natural conjecture in the field.

2. PRELIMINARIES

2.1. Function spaces. For $p \geq 1$ and measurable $f : \mathbb{R}^n \rightarrow \mathbb{R}$, let

$$\|f\|_p = \left(\int_{\mathbb{R}^n} |f(x)|^p dx \right)^{1/p}.$$

We set $\{f \geq t\} = \{x \in \mathbb{R}^n : f(x) \geq t\}$ for $t \in \mathbb{R}$ and use corresponding notation for level sets, etc. We say that f is non-zero, if $\{f \neq 0\}$ has positive measure, and we identify functions that are equal up to a set of measure zero. For $p \geq 1$, let

$$L^p(\mathbb{R}^n) = \left\{ f : \mathbb{R}^n \rightarrow \mathbb{R} : f \text{ is measurable, } \|f\|_p < \infty \right\}.$$

Here and below, when we use measurability and related notions, we refer to the n -dimensional Lebesgue measure on \mathbb{R}^n .

For $E \subset \mathbb{R}^n$, the indicator function 1_E is defined by $1_E(x) = 1$ for $x \in E$ and $1_E(x) = 0$ otherwise. We say that two measurable sets are equivalent if their indicator functions are equal up to a set of measure zero.

Let $0 < s < 1$. We define the s -fractional Sobolev space as

$$W^{s,1}(\mathbb{R}^n) = \left\{ f \in L^1(\mathbb{R}^n) : \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy < \infty \right\}.$$

For $f \in L^1(\mathbb{R}^n)$, we say that f admits a weak gradient $\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ if

$$\int_{\mathbb{R}^n} \langle \nu(x), \nabla f(x) \rangle dx = - \int_{\mathbb{R}^n} f(x) \operatorname{div} \nu(x) dx$$

for every $\nu \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$, where $C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$ is the set of smooth vector fields $\nu : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with compact support and $\operatorname{div} \nu$ denotes the divergence of ν . We set

$$W^{1,1}(\mathbb{R}^n) = \left\{ f \in L^1(\mathbb{R}^n) : |\nabla f| \in L^1(\mathbb{R}^n) \right\}.$$

Let $\mathcal{B}(\mathbb{R}^n)$ denote the class of Borel sets in \mathbb{R}^n . For $f \in L^1(\mathbb{R}^n)$, we say that f is a function of bounded variation on \mathbb{R}^n if there is a finite vector-valued Radon measure $Df : \mathcal{B}(\mathbb{R}^n) \rightarrow \mathbb{R}^n$ such that

$$(8) \quad \int_{\mathbb{R}^n} \langle \nu(x), \sigma_f(x) \rangle d|Df|(x) = - \int_{\mathbb{R}^n} f(x) \operatorname{div} \nu(x) dx$$

for every $\nu \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$, where $|Df| : \mathcal{B}(\mathbb{R}^n) \rightarrow [0, \infty)$ denotes the variation measure of Df and $\sigma_f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ the Radon–Nikodym derivative of Df with respect to $|Df|$. We write $BV(\mathbb{R}^n)$ for the space of L^1 functions of bounded variation. For more information on functions of bounded variation, we refer to [8, Chapter 5].

2.2. Symmetrization. Let $E \subseteq \mathbb{R}^n$ be a Borel set of finite measure. The Schwarz symmetral of E , denoted by E^* , is the centered Euclidean ball with same volume as E .

For a non-negative integrable function f and $t > 0$, the superlevel set $\{f \geq t\}$ is a measurable set of finite measure. The layer cake formula states that

$$f(x) = \int_0^\infty 1_{\{f \geq t\}}(x) dt$$

for almost every $x \in \mathbb{R}^n$ and allows us to recover the function from its superlevel sets. The Schwarz symmetral of f , denoted by f^* , is defined by

$$f^*(x) = \int_0^\infty 1_{\{f \geq t\}^*}(x) dt$$

for $x \in \mathbb{R}^n$. Hence f^* is determined by the properties of being radially symmetric and having superlevel sets that are balls of the same measure as the superlevel sets of f . Note that f^* is often called symmetric decreasing rearrangement of f .

Let $\xi \in S^{n-1}$. The Steiner symmetral of E in the direction ξ , denoted by E^ξ , is defined by the property that for every line L parallel to ξ , the set $L \cap E^\xi$ is an interval of the same length as $L \cap E$ and symmetric with respect to the subspace orthogonal to ξ .

For a non-negative integrable function f , the Steiner symmetral f^ξ is defined by

$$f^\xi(x) = \int_0^\infty 1_{\{f \geq t\}^\xi}(x) dt$$

for $x \in \mathbb{R}^n$.

2.3. Star bodies. For a set $K \subseteq \mathbb{R}^n$ that is star-shaped (with respect to the origin), the gauge function $\|\cdot\|_K : \mathbb{R}^n \rightarrow [0, \infty]$ is defined as

$$\|x\|_K = \inf\{\lambda > 0 : x \in \lambda K\},$$

and the radial function $\rho_K : \mathbb{R}^n \setminus \{0\} \rightarrow [0, \infty]$ as

$$\rho_K(x) = \|x\|_K^{-1} = \sup\{\lambda \geq 0 : \lambda x \in K\}.$$

We call K a star body if its radial function is strictly positive and continuous in $\mathbb{R}^n \setminus \{0\}$.

We say that a star-shaped set is p -convex with $0 < p \leq 1$ if

$$\|x + y\|_K^p \leq \|x\|_K^p + \|y\|_K^p$$

for all $x, y \in \mathbb{R}^n$. A p -convex body is a star body that is also p -convex. For more information on p -convex sets, see, for example, [16].

For star bodies $K, L \subseteq \mathbb{R}^n$ and $p \in \mathbb{R}$, define the dual mixed volume as

$$\tilde{V}_p(K, L) = \frac{1}{n} \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n-p} \rho_L(\xi)^p d\xi.$$

Notice that

$$\tilde{V}_p(K, K) = |K|.$$

The dual mixed volume inequality (see [31, Section 9.3] or [12, B.29]) states that for $p < 0$ or $p > n$,

$$(9) \quad \tilde{V}_p(K, L) \geq |K|^{(n-p)/n} |L|^{p/n},$$

and the reverse inequality holds for $0 < p < n$. Equality holds for $p \neq 0, n$ if and only if K and L are dilates, where we say that star bodies K and L are dilates if $\rho_K = c \rho_L$ on \mathbb{S}^{n-1} for some $c > 0$.

2.4. Convex bodies and the Petty projection inequality. A set $K \subseteq \mathbb{R}^n$ is called a (proper) convex body if it is compact and convex and has non-empty interior. For a convex body K , the support function $h_K : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as

$$h_K(y) = \max\{\langle x, y \rangle : x \in K\}.$$

A convex body K that contains the origin in its interior is also a star body. The polar body of such K is defined as

$$K^* = \{y \in \mathbb{R}^n : \langle x, y \rangle \leq 1 \text{ for all } x \in K\}$$

and is a convex body containing the origin in its interior while

$$\|x\|_K = h_{K^*}(x)$$

for $x \in \mathbb{R}^n$.

For $K \subset \mathbb{R}^n$ a convex body, the projection body is defined as the convex body whose support function for $\xi \in \mathbb{S}^{n-1}$ is

$$h_{\Pi K}(\xi) = |\text{proj}_{\xi^\perp}(K)|_{n-1},$$

where proj_{ξ^\perp} denotes the orthogonal projection to the hyperplane, ξ^\perp , orthogonal to ξ and $|\cdot|_{n-1}$ stands for $(n-1)$ -dimensional volume in ξ^\perp . Using the surface area measure $S(K, \cdot)$ of K , we can also write

$$h_{\Pi K}(\xi) = \frac{1}{2} \int_{\mathbb{S}^{n-1}} |\langle \xi, \eta \rangle| dS(K, \eta)$$

for $\xi \in \mathbb{S}^{n-1}$. Projection bodies were introduced by Minkowski and $h_{\Pi K}(\xi)$ is called the brightness of K in the direction ξ (see [12, 31] for more information).

The important Petty projection inequality [28] states that for a convex body $K \subset \mathbb{R}^n$,

$$(10) \quad |K|^{(n-1)/n} \leq \frac{\omega_n}{\omega_{n-1}} |\Pi^* K|^{-1/n}$$

with equality precisely for ellipsoids, where $\Pi^* K$ is the polar body of ΠK . A new proof using Theorem 1 (and hence the Riesz rearrangement inequality) is given in Section 7. A previous proof of (10), that also uses the Riesz rearrangement inequality and the so-called convolution square of a convex body, was given by Schmuckenschläger [32]. For generalization of projection bodies of convex bodies, see, for example, [2, 6, 14, 18, 20, 23, 26], and for further results related to the Sobolev–Zhang inequality, see, for example, [15, 24, 25, 27, 36].

2.5. Anisotropic fractional Sobolev norms and perimeters. Let $0 < s < 1$ and let $K \subset \mathbb{R}^n$ be a star body. For $f \in W^{s,1}(\mathbb{R}^n)$, the anisotropic fractional Sobolev norm of f with respect to K is

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K^{n+s}} dx dy.$$

It was introduced in [21] for K a convex body (also, see [22]). The s -fractional perimeter with respect to K of a measurable set $E \subseteq \mathbb{R}^n$ was introduced in [21] as

$$P_s(E, K) = \int_E \int_{E^c} \frac{1}{\|x - y\|_K^{n+s}} dx dy,$$

where $E^c = \mathbb{R}^n \setminus E$ is the complement of E in \mathbb{R}^n . The same definition will be used for $K \subset \mathbb{R}^n$ a star body. The anisotropic s -fractional perimeter of E with respect to K is half of the s -fractional anisotropic Sobolev norm of 1_E with respect to K . Since K is a star body, this implies for measurable $E \subset \mathbb{R}^n$ that $P_s(E, K) < \infty$ if and only if $1_E \in W^{s,1}(\mathbb{R}^n)$. If K is the Euclidean unit ball, we write $P_s(E)$ and obtain the well-known Euclidean s -fractional perimeter of E . We will use that

$$(11) \quad P_s(B^n) = \frac{2^{1-s} \pi^{(n-1)/2} n \omega_n \Gamma\left(\frac{1-s}{2}\right)}{s(n-s) \Gamma\left(\frac{n-s}{2}\right)} = \frac{2^{1-s} n \omega_n \omega_{n-s}}{s(1-s) \omega_{1-s}}$$

(cf. [9]), where $\omega_q = \pi^{q/2} / \Gamma(q/2 + 1)$ for $q > 0$ and Γ is the gamma function. A simple way to obtain this result is by using the Blaschke–Petkantschin formula [33, Theorem 7.2.7]:

$$\begin{aligned} P_s(B^n) &= \int_{B^n \cap L \neq \emptyset} \int_{B^n \cap L} \int_{(\mathbb{R}^n \setminus B^n) \cap L} \frac{dH^1(x) dH^1(y)}{|x - y|^{s+1}} dL \\ &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} \int_{\xi^\perp} \int_{B^n \cap (z + \mathbb{R}\xi)} \int_{(\mathbb{R}^n \setminus B^n) \cap (z + \mathbb{R}\xi)} \frac{dH^1(x) dH^1(y)}{|x - y|^{1+s}} dz d\xi \\ &= \frac{n \omega_n (n-1) \omega_{n-1}}{2} \int_0^1 \int_{|x| \leq \sqrt{1-t^2}} \int_{|y| \geq \sqrt{1-t^2}} \frac{dH^1(x) dH^1(y)}{|x - y|^{1+s}} t^{n-2} dt \\ &= \frac{2^{1-s} (n-1) n \omega_{n-1} \omega_n}{s(1-s)} \int_0^1 (1-t^2)^{(1-s)/2} t^{n-2} dt \\ &= \frac{(n-1) n \omega_{n-1} \omega_n}{2^s s(1-s)} B\left(\frac{1-s}{2} + 1, \frac{n-1}{2}\right) \end{aligned}$$

where the outer integration in the first integral is on the affine Grassmannian of lines in \mathbb{R}^n and we use the $(n-1)$ -dimensional Hausdorff measure for the direction of lines and the $(n-1)$ -dimensional Lebesgue measure in the orthogonal complement of a line. Here H^1 denotes 1-dimensional Hausdorff measure and $\mathbb{R}\xi$ is the line with direction ξ while B is the beta function. For the optimal constant in (1), we get

$$\alpha_{n,s} = 2P_s(B^n) \omega_n^{-(n-s)/n}$$

since there is equality in (1) for indicator functions of balls.

The following fractional co-area formula is a consequence of Fubini's theorem (cf. [34, Example (2.9)]):

$$(12) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K^{n+s}} dx dy = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \int_0^\infty \frac{|1_{\{f \geq t\}}(x) - 1_{\{f \geq t\}}(y)|}{\|x - y\|_K^{n+s}} dt dx dy$$

$$= 2 \int_0^\infty P_s(\{f \geq t\}, K) dt$$

for non-negative $f \in W^{s,1}(\mathbb{R}^n)$ and a star body $K \subset \mathbb{R}^n$.

3. FRACTIONAL POLAR PROJECTION BODIES

Let $0 < s < 1$ and $f \in BV(\mathbb{R}^n)$. We discuss properties of the s -fractional polar projection body $\Pi^{*,s}f$, that is, the star-shaped set defined through its gauge function for $\xi \in \mathbb{R}^n$ as

$$(13) \quad \|\xi\|_{\Pi^{*,s}f}^s = \int_0^\infty t^{-s} \int_{\mathbb{R}^n} \left| \frac{f(x + t\xi) - f(x)}{t} \right| dx dt.$$

Notice that $\|\cdot\|_{\Pi^{*,s}f}$ is a 1-homogeneous function.

We start with a simple observation. Let $K \subset \mathbb{R}^n$ be a star body. Using (13), we obtain

$$\begin{aligned} & \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K^{n+s}} dx dy \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(y + z) - f(y)|}{\|z\|_K^{n+s}} dz dy \\ &= \int_{\mathbb{S}^{n-1}} \int_0^\infty \|t\xi\|_K^{-n-s} \int_{\mathbb{R}^n} |f(y + t\xi) - f(y)| t^{n-1} dy dt d\xi \\ &= \int_{\mathbb{S}^{n-1}} \|\xi\|_K^{-n-s} \int_0^\infty t^{-s} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 dt d\xi \\ &= \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n+s} \rho_{\Pi^{*,s}f}(\xi)^{-s} d\xi. \end{aligned}$$

Hence

$$(14) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K^{n+s}} dx dy = n \tilde{V}_{-s}(K, \Pi^{*,s}f),$$

where we have already used the following result, which shows that $\Pi^{*,s}f$ is an origin-symmetric star body.

Proposition 2. *For non-zero $f \in W^{s,1}(\mathbb{R}^n)$, the set $\Pi^{*,s}f$ is an origin-symmetric s -convex body with the origin in its interior. Moreover, there is $c > 0$ depending only on f such that $\Pi^{*,s}f \subseteq cB^n$ for every $s \in (0, 1)$.*

Proof. First, note that since for $\xi \in \mathbb{R}^n$ and $t > 0$,

$$\int_{\mathbb{R}^n} |f(x - t\xi) - f(x)| dx = \int_{\mathbb{R}^n} |f(x) - f(x + t\xi)| dx,$$

the set $\Pi^{*,s}f$ is origin-symmetric.

Next, we show that $\Pi^{*,s}f$ is bounded. We take $r > 1$ large enough so that $\int_{rB^n} |f(x)| dx \geq \frac{2}{3}\|f\|_1$ and easily see that for $t > r$,

$$\begin{aligned} & \int_{\mathbb{R}^n} |f(x+t\xi) - f(x)| dx \\ & \geq \int_{rB^n} (|f(x)| - |f(x+t\xi)|) dx + \int_{rB^n - t\xi} (|f(x+t\xi)| - |f(x)|) dx \\ & \geq 2\left(\frac{2}{3}\|f\|_1 - \frac{1}{3}\|f\|_1\right). \end{aligned}$$

Hence,

$$\int_0^\infty t^{-s} \left\| \frac{f(x+t\xi) - f(x)}{t} \right\|_1 dt \geq \frac{2}{3}\|f\|_1 \int_r^\infty t^{-s-1} dt \geq \frac{2}{3r}\|f\|_1,$$

which implies that $\Pi^{*,s}f \subseteq cB^n$ for $c > 0$ independent of s .

Next, we show that $\|\cdot\|_{\Pi^{*,s}f}^s$ is sublinear on \mathbb{R}^n . Indeed, for $\xi, \eta \in \mathbb{R}^n$, the triangle inequality and a change of variables show that

$$\begin{aligned} (15) \quad \|\xi + \eta\|_{\Pi^{*,s}f}^s &= \int_0^\infty t^{-s-1} \|f(\cdot + t\xi + t\eta) - f(\cdot)\|_1 dt \\ &\leq \int_0^\infty t^{-s-1} (\|f(\cdot + t\xi + t\eta) - f(\cdot + t\xi)\|_1 + \|f(\cdot + t\xi) - f(\cdot)\|_1) dt \\ &= \|\xi\|_{\Pi^{*,s}f}^s + \|\eta\|_{\Pi^{*,s}f}^s, \end{aligned}$$

which shows that $\Pi^{*,s}f$ is an s -convex set.

Now, we show that $\Pi^{*,s}f$ has the origin in its interior. Using the relation (14), we get

$$\int_{\mathbb{S}^{n-1}} \|\xi\|_{\Pi^{*,s}f}^s d\xi = \frac{1}{n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy,$$

which is finite since $f \in W^{s,1}(\mathbb{R}^n)$. We choose $r > 0$ large enough so that the set $A = \{\xi \in \mathbb{S}^{n-1} : \|\xi\|_{\Pi^{*,s}f}^s < r\}$ has positive $(n-1)$ -dimensional Hausdorff measure and a basis $\{\xi_1, \dots, \xi_n\} \subseteq A$ of \mathbb{R}^n . Applying (if necessary) a linear transformation to $\Pi^{*,s}f$, we may assume without loss of generality that $\xi_i = e_i$ are the canonical basis vectors. For every $x \in \mathbb{R}^n$ writing $x = \sum x_i e_i$ and using (15) we get

$$\|x\|_{\Pi^{*,s}f} \leq \left(\sum |x_i|^s \|e_i\|_{\Pi^{*,s}f}^s \right)^{1/s} \leq d|x|,$$

where $d > 0$ is independent of x . This shows that $\Pi^{*,s}f$ has the origin as interior point.

Finally, it is easy to see that a star-shaped set that is s -convex and has the origin in its interior, also has a continuous gauge function, being thus an s -convex body. \square

4. THE LIMIT CASE $s \rightarrow 1^-$ OF FRACTIONAL POLAR PROJECTION BODIES

We prove (6) and the corresponding statement for functions of bounded variation. For $f \in BV(\mathbb{R}^n)$, the polar projection body is defined for $\xi \in \mathbb{S}^{n-1}$ by

$$\|\xi\|_{\Pi^*f} = \frac{1}{2} \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x)$$

(see [35]). Note that for $f \in W^{1,1}(\mathbb{R}^n)$, this definition coincides with (4).

Theorem 3. Let $f \in BV(\mathbb{R}^n)$. For $\xi \in \mathbb{S}^{n-1}$,

$$\lim_{s \rightarrow 1^-} (1-s) \|\xi\|_{\Pi^{*,s}f}^s = 2 \|\xi\|_{\Pi^*f}.$$

Moreover,

$$\lim_{s \rightarrow 1^-} (1-s) |\Pi^{*,s}f|^{-s/n} = 2 |\Pi^*f|^{-1/n}$$

and

$$\lim_{s \rightarrow 1^-} (1-s) \tilde{V}_{-s}(K, \Pi^{*,s}f) = 2 \tilde{V}_{-1}(K, \Pi^*f)$$

for every star body $K \subset \mathbb{R}^n$.

We require the following two lemmas.

Lemma 4. If $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a measurable function with $\lim_{t \rightarrow 0^+} \varphi(t) = \varphi(0)$ and such that $\int_0^\infty t^{-s_0} \varphi(t) dt < \infty$ for some $s_0 \in (0, 1)$, then

$$\lim_{s \rightarrow 1^-} (1-s) \int_0^\infty t^{-s} \varphi(t) dt = \varphi(0).$$

Proof. Given $\varepsilon > 0$ take $\delta \in (0, 1)$ so that $0 \leq t \leq \delta$ implies $|\varphi(t) - \varphi(0)| \leq \varepsilon$. We have

$$\begin{aligned} \int_0^\infty t^{-s} \varphi(t) dt &= \int_0^\delta t^{-s} \varphi(0) dt + \int_0^\delta t^{-s} (\varphi(t) - \varphi(0)) dt + \int_\delta^\infty t^{-s} \varphi(t) dt \\ &= \frac{\delta^{1-s}}{1-s} \varphi(0) + A + B \end{aligned}$$

where $(1-s)|A| \leq \varepsilon \delta^{1-s} \leq \varepsilon$ and

$$|B| \leq \delta^{s_0-s} \int_\delta^\infty t^{-s_0} \varphi(t) dt \leq \delta^{s_0-s} \int_0^\infty t^{-s_0} \varphi(t) dt$$

for $s \in (s_0, 1)$. Since $(1-s)B \rightarrow 0$ and $\delta^{1-s} \rightarrow 1$ as $s \rightarrow 1^-$, we can take s sufficiently close to 1 so that $(1-s)|B| \leq \varepsilon$ and $|\delta^{1-s} \varphi(0) - \varphi(0)| \leq \varepsilon$, obtaining $|(1-s) \int_0^\infty t^{-s} \varphi(t) dt - \varphi(0)| \leq 3\varepsilon$. \square

Lemma 5. For $f \in BV(\mathbb{R}^n)$ and $\xi \in \mathbb{S}^{n-1}$,

$$\lim_{t \rightarrow 0} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 = \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x).$$

Proof. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function with compact support, and write div_x for the divergence taken with respect to the variable x . Using (8), we obtain

$$\begin{aligned} \int_{\mathbb{R}^n} g(x) \frac{f(x + t\xi) - f(x)}{t} dx &= \int_{\mathbb{R}^n} f(x) \frac{g(x - t\xi) - g(x)}{t} dx \\ &= - \int_{\mathbb{R}^n} f(x) \int_0^1 \langle \nabla g(x - rt\xi), \xi \rangle dr dx \\ &= - \int_{\mathbb{R}^n} f(x) \operatorname{div}_x \left(\int_0^1 g(x - rt\xi) dr \xi \right) dx \\ &= \int_{\mathbb{R}^n} \left(\int_0^1 g(x - rt\xi) dr \right) \langle \sigma_f(x), \xi \rangle d|Df|(x). \end{aligned}$$

Therefore

$$(16) \quad \lim_{t \rightarrow 0} \int_{\mathbb{R}^n} g(x) \frac{f(x+t\xi) - f(x)}{t} dx = \int_{\mathbb{R}^n} g(x) \langle \sigma_f(x), \xi \rangle d|Df|(x).$$

Let $\varepsilon > 0$. Since $|Df|$ is a finite Radon measure, the set of smooth functions with compact support is dense in the space of L^1 functions w.r.t. $|Df|$ (cf. [10, Proposition 7.9]). Hence, there is a smooth function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ with compact support such that

$$\int_{\mathbb{R}^n} |g(x) - \operatorname{sgn}(\langle \sigma_f(x), \xi \rangle)| d|Df|(x) < \varepsilon$$

and $\|g\|_\infty \leq 1 + \varepsilon$. By (16),

$$\begin{aligned} (1 + \varepsilon) \liminf_{t \rightarrow 0} \int_{\mathbb{R}^n} \left| \frac{f(x+t\xi) - f(x)}{t} \right| dx &\geq \lim_{t \rightarrow 0} \int_{\mathbb{R}^n} g(x) \frac{f(x+t\xi) - f(x)}{t} dx \\ &= \int_{\mathbb{R}^n} g(x) \langle \sigma_f(x), \xi \rangle d|Df|(x) \\ &= \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x) \\ &\quad + \int_{\mathbb{R}^n} (g(x) - \operatorname{sgn}(\langle \sigma_f(x), \xi \rangle)) \langle \sigma_f(x), \xi \rangle d|Df|(x) \\ &\geq \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x) - \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ was arbitrary, we obtain

$$\liminf_{t \rightarrow 0} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 \geq \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x).$$

For the opposite inequality, notice that

$$\left| \int_{\mathbb{R}^n} \left(\int_0^1 h(x - rt\xi) dr \right) \langle \sigma_f(x), \xi \rangle d|Df|(x) \right| \leq \|h\|_\infty \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x)$$

for every $h \in L^\infty(\mathbb{R}^n)$. □

Proof of Theorem 3. Define $\varphi : [0, \infty) \rightarrow [0, \infty)$ by

$$\varphi(t) = \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1$$

and note that $\varphi(t) \leq \frac{2\|f\|_1}{t}$ for $t > 0$. By Lemma 4 and Lemma 5,

$$\lim_{s \rightarrow 1^-} (1-s) \int_0^\infty t^{-s} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 dt = \int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x).$$

By Proposition 2 we can use the dominated convergence theorem to obtain,

$$\begin{aligned}
 & \lim_{s \rightarrow 1^-} n |(1-s)^{-1/s} \Pi^{*,s} f| \\
 &= \lim_{s \rightarrow 1^-} \int_{\mathbb{S}^{n-1}} \left((1-s) \int_0^\infty t^{-s} \left\| \frac{f(\cdot + t\xi) - f(\cdot)}{t} \right\|_1 dt \right)^{-n/s} d\xi \\
 &= \int_{\mathbb{S}^{n-1}} \left(\int_{\mathbb{R}^n} |\langle \sigma_f(x), \xi \rangle| d|Df|(x) \right)^{-n} d\xi \\
 &= n |\frac{1}{2} \Pi^* f|
 \end{aligned}$$

and

$$\begin{aligned}
 \lim_{s \rightarrow 1^-} n(1-s) \tilde{V}_{-s}(K, \Pi^{*,s} f) &= \lim_{s \rightarrow 1^-} (1-s) \int_{\mathbb{S}^{n-1}} \|\xi\|_K^{n+s} \|\xi\|_{\Pi^{*,s} f}^s d\xi \\
 &= 2 \int_{\mathbb{S}^{n-1}} \|\xi\|_K^n \|\xi\|_{\Pi^* f} d\xi \\
 &= 2n \tilde{V}_{-1}(K, \Pi^* f),
 \end{aligned}$$

which completes the proof of the theorem. \square

5. ANISOTROPIC FRACTIONAL PÓLYA–SZEGŐ INEQUALITIES

Almgren and Lieb [1] established the following Pólya–Szegő inequality for fractional Sobolev norms. The equality case was settled by Frank and Seiringer [11]. Let $0 < s < 1$ and $n \geq 1$. For non-negative $f \in W^{s,1}(\mathbb{R}^n)$,

$$(17) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy \geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f^*(x) - f^*(y)|}{|x - y|^{n+s}} dx dy,$$

with equality precisely if $\{f \geq t\}$ is equivalent to a ball for almost every $t > 0$.

The following improvement of (17) is key to prove the main theorems in Section 8. It is a variation of [17, Theorem 3.1].

Theorem 6. *Let $0 < s < 1$ and $K \subset \mathbb{R}^n$ a star body. For $f : \mathbb{R}^n \rightarrow \mathbb{R}$ a non-negative integrable function,*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K^{n+s}} dx dy \geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f^*(x) - f^*(y)|}{\|x - y\|_{K^*}^{n+s}} dx dy.$$

If equality holds for non-zero $f \in W^{s,1}(\mathbb{R}^n)$, then K is a centered ellipsoid and for almost every $t > 0$, the level set $\{f \geq t\}$ is equivalent to an ellipsoid.

The inequality in Theorem 6 follows rather directly from the Riesz rearrangement inequality, which is stated in full generality, for example, in [4].

Theorem 7 (Riesz's rearrangement inequality). *For $f, g, k : \mathbb{R}^n \rightarrow \mathbb{R}$ non-negative measurable functions,*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x) k(x - y) g(y) dx dy \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f^*(x) k^*(x - y) g^*(y) dx dy.$$

To establish the equality case in Theorem 6, we use the characterization of equality cases in a version of the Riesz rearrangement inequality due to Burchard [5].

Theorem 8 (Burchard). *Let A, B and C be sets of finite positive measure in \mathbb{R}^n and denote by α, β and γ the radii of their Schwarz symmetrals A^*, B^* and C^* . For $|\alpha - \beta| < \gamma < \alpha + \beta$, there is equality in*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_A(y) 1_B(x-y) 1_C(x) dx dy \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_{A^*}(y) 1_{B^*}(x-y) 1_{C^*}(x) dx dy$$

if and only if, up to sets of measure zero,

$$A = a + \alpha D, B = b + \beta D, C = c + \gamma D,$$

where D is a centered ellipsoid, and a, b and $c = a + b$ are vectors in \mathbb{R}^n .

By the co-area formula (12), Theorem 6 is an immediate consequence of the following result for anisotropic fractional perimeters.

Theorem 9. *Let $0 < s < 1$ and $K \subset \mathbb{R}^n$ a star body. For $E \subset \mathbb{R}^n$ measurable,*

$$(18) \quad \int_E \int_{E^c} \frac{1}{\|x-y\|_K^{n+s}} dx dy \geq \int_{E^*} \int_{(E^*)^c} \frac{1}{\|x-y\|_{K^*}^{n+s}} dx dy.$$

If equality holds for E with finite positive s -perimeter, then K is a centered ellipsoid and E is equivalent to an ellipsoid.

Proof. Write

$$\|z\|_K^{-n-s} = \int_0^\infty k_t(z) dt, \text{ where } k_t(z) = 1_{t^{-1/(n+s)}K}(z)$$

and use Fubini's theorem, to obtain

$$\int_E \int_{E^c} \frac{1}{\|x-y\|_K^{n+s}} dx dy = \int_0^\infty \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_E(x) 1_{E^c}(y) k_t(x-y) dx dy dt.$$

For fixed $t \in (0, \infty)$, we have

$$(19) \quad \begin{aligned} & \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_E(x) 1_{E^c}(y) k_t(x-y) dx dy \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} (1_E(x) - 1_E(x) 1_E(y)) k_t(x-y) dx dy \\ &= t^{-\frac{n}{n+s}} |K| |E| - \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_E(x) k_t(x-y) 1_E(y) dx dy. \end{aligned}$$

Clearly the first term is invariant under Schwarz symmetrization, and for the second term we can apply Theorem 7 to show that the left side does not increase under Schwarz symmetrization. Thus

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_E(x) k_t(x-y) 1_{E^c}(y) dx dy \geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_{E^*}(x) k_t^*(x-y) 1_{(E^*)^c}(y) dx dy$$

for $t > 0$ and integrating this inequality for $t \in (0, \infty)$, we obtain (18).

If there is equality in (18), it follows from (19) that for almost every $t > 0$,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} 1_E(x) 1_{t^{-1/(n+s)}K}(x-y) 1_E(y) dx dy$$

is invariant under Schwarz symmetrization. Now we may apply Theorem 8 with $A = C = E$ and $B = t^{-1/(n+s)}K$ for suitable $t > 0$. Since $A = C$ we have $a = c$ and this implies $b = 0$. Hence K is a centered ellipsoid and E is equivalent to an ellipsoid. \square

To establish corresponding inequalities for Steiner symmetrization, we require the following result.

Theorem 10 (Rogers [29]). *Let $\xi \in \mathbb{S}^{n-1}$. For $f, g, k : \mathbb{R}^n \rightarrow \mathbb{R}$ non-negative and measurable,*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x)k(x-y)g(y) dx dy \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f^\xi(x)k^\xi(x-y)g^\xi(y) dx dy.$$

We remark that the above theorem is a special case of the inequalities of Rogers [30] and Brascamp–Lieb–Luttinger [4, Lemma 3.2].

As in the case of Schwarz symmetrization, we obtain the following consequences.

Theorem 11. *Let $0 < s < 1$ and $\xi \in \mathbb{S}^{n-1}$. If $K \subset \mathbb{R}^n$ is a star body, then*

$$\int_E \int_{E^c} \frac{1}{\|x-y\|_K^{n+s}} dx dy \geq \int_{E^\xi} \int_{(E^\xi)^c} \frac{1}{\|x-y\|_{K^\xi}^{n+s}} dx dy$$

for $E \subset \mathbb{R}^n$ measurable.

Theorem 12. *Let $0 < s < 1$ and $\xi \in \mathbb{S}^{n-1}$. If $K \subset \mathbb{R}^n$ is a star body, then*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x-y\|_K^{n+s}} dx dy \geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f^\xi(x) - f^\xi(y)|}{\|x-y\|_{K^\xi}^{n+s}} dx dy$$

for $f : \mathbb{R}^n \rightarrow \mathbb{R}$ non-negative and integrable.

6. AFFINE FRACTIONAL PÓLYA–SZEGŐ INEQUALITIES

We establish the following affine version of the fractional Pólya–Szegő inequality (17) by Almgren and Lieb.

Theorem 13. *For $0 < s < 1$ and non-negative $f \in W^{s,1}(\mathbb{R}^n)$,*

$$(20) \quad |\Pi^{*,s} f|^{-s/n} \geq |\Pi^{*,s} f^*|^{-s/n},$$

with equality precisely if $\{f \geq t\}$ is equivalent to an ellipsoid for almost every $t > 0$.

Proof. By Theorem 6, (14) and (9), we have

$$\begin{aligned} \tilde{V}_{-s}(K, \Pi^{*,s} f) &\geq \tilde{V}_{-s}(K^*, \Pi^{*,s} f^*) \\ &\geq |K^*|^{(n+s)/n} |\Pi^{*,s} f^*|^{-s/n} \\ &= |K|^{(n+s)/n} |\Pi^{*,s} f^*|^{-s/n}. \end{aligned}$$

Setting $K = \Pi^{*,s} f$, we obtain that

$$|\Pi^{*,s} f| = \tilde{V}_{-s}(\Pi^{*,s} f, \Pi^{*,s} f) \geq |\Pi^{*,s} f|^{(n+s)/n} |\Pi^{*,s} f^*|^{-\frac{s}{n}},$$

which completes the proof of the inequality. The equality case follows from the equality case of the dual mixed volumes inequality and of Theorem 6. \square

If we multiply (20) by $(1-s)$ and let $s \rightarrow 1^-$, we obtain from Theorem 13 and Theorem 3 the following affine Pólya–Szegő inequality by Cianchi, Lutwak, Yang and Zhang [6] (see also [27]):

$$|\Pi^* f|^{-1/n} \geq |\Pi^* f^*|^{-1/n}$$

for $f \in W^{1,1}(\mathbb{R}^n)$.

Using the same proof as for Theorem 13 for Steiner symmetrization and replacing Theorem 7 by Theorem 12, we obtain the following result.

Theorem 14. *Let $0 < s < 1$ and $\xi \in \mathbb{S}^{n-1}$. Then*

$$|\Pi^{*,s} f|^{-s/n} \geq |\Pi^{*,s} f^\xi|^{-s/n}$$

for non-negative $f \in W^{s,1}(\mathbb{R}^n)$.

As before, we obtain from Theorem 14 and Theorem 3 the following affine inequality:

$$|\Pi^* f|^{-1/n} \geq |\Pi^* f^\xi|^{-1/n}$$

for $f \in W^{1,1}(\mathbb{R}^n)$ and $\xi \in \mathbb{S}^{n-1}$.

7. FRACTIONAL PETTY PROJECTION INEQUALITIES

A set $E \subset \mathbb{R}^n$ of finite measure is a set of finite perimeter if its indicator function 1_E is in $BV(\mathbb{R}^n)$. This allows us to translate results for functions in $BV(\mathbb{R}^n)$ to sets of finite perimeter and finite measure. The following result is an immediate consequence of the affine fractional Pólya–Szegő inequality from Theorem 13.

Theorem 15. *For $0 < s < 1$ and $E \subset \mathbb{R}^n$ of finite s -perimeter and finite measure,*

$$|\Pi^{*,s} E|^{-s/n} \geq |\Pi^{*,s} E^*|^{-s/n}$$

with equality precisely if E is equivalent to an ellipsoid.

Here we write $\Pi^{*,s} E$ for the fractional polar projection body of 1_E . Note that

$$\|\xi\|_{\Pi^{*,s} E}^s = \int_0^\infty t^{-s} \frac{|E \Delta (E - t\xi)|}{t} dt$$

for $E \subset \mathbb{R}^n$ of finite s -perimeter and finite measure. Also note that (14) implies that

$$(21) \quad 2 \int_E \int_{E^c} \frac{1}{\|x-y\|_K^{n+s}} dx dy = n \tilde{V}_{-s}(K, \Pi^{*,s} E)$$

for $K \subset \mathbb{R}^n$ a star body.

From Theorem 15, we easily obtain the first inequality of the following result, which we call the fractional Petty projection inequality.

Theorem 16. *For $0 < s < 1$ and $E \subset \mathbb{R}^n$ of finite s -perimeter and finite measure,*

$$(22) \quad \left(\frac{|E|}{|B^n|} \right)^{(n-s)/n} \leq \left(\frac{|\Pi^{*,s} E|}{|\Pi^{*,s} B^n|} \right)^{-s/n} \leq \frac{P_s(E)}{P_s(B^n)}.$$

There is equality in the first inequality if and only if E is equivalent to an ellipsoid. There is equality in the second inequality if and only if E is equivalent to a set of constant s -fractional brightness.

Here, we say that $E \subset \mathbb{R}^n$ is of constant s -fractional brightness if $\Pi^{*,s}E$ is a dilate of B^n . The second inequality in (22) and its equality case follow directly from (14) and the dual mixed volume inequality (9).

Let $E \subset \mathbb{R}^n$ be a set of finite perimeter and finite measure. The perimeter of E is defined as $P(E) = |D 1_E|$. By Dávila's version of (2),

$$\lim_{s \rightarrow 1^-} \frac{P_s(E)}{P_s(B^n)} = \frac{P(E)}{P(B^n)}.$$

Taking the limit $s \rightarrow 1^-$ in (22), we obtain from this and Theorem 3 that

$$(23) \quad \left(\frac{|E|}{|B^n|}\right)^{(n-1)/n} \leq \left(\frac{|\Pi^*E|}{|\Pi^*B^n|}\right)^{-1/n} \leq \frac{P(E)}{P(B^n)}$$

for $E \subset \mathbb{R}^n$ of finite perimeter and finite measure. The first inequality is the generalized Petty projection inequality that was proved by Gaoyong Zhang [38] for compact sets with piecewise C^1 boundary and in full generality by Tuo Wang [35]. For E a convex body, it is the classical Petty projection inequality (10). There is equality in the second inequality of (23) if and only if E is of constant brightness.

The Steiner inequality for fractional polar projection bodies is contained in the following result, which is an immediate consequence of Theorem 14.

Theorem 17. *Let $0 < s < 1$ and $\xi \in \mathbb{S}^{n-1}$. Then*

$$|\Pi^{*,s}E|^{-s/n} \geq |\Pi^{*,s}E^\xi|^{-s/n}$$

for $E \subset \mathbb{R}^n$ of finite s -perimeter and finite Lebesgue measure.

Taking the limit as $s \rightarrow 1^-$ and using Theorem 3, we obtain from Theorem 17 the following Steiner inequality for polar projection bodies, which was recently established by Youjiang Lin [19]:

$$|\Pi^*E|^{-1/n} \geq |\Pi^*E^\xi|^{-1/n}$$

for $E \subset \mathbb{R}^n$ of finite perimeter and finite Lebesgue measure and $\xi \in \mathbb{S}^{n-1}$.

8. PROOF OF THEOREM 1

A version of the following lemma for $s = 1$ is contained in [38].

Lemma 18. *Let $0 < s < 1$. If $g : \mathbb{R}^n \rightarrow [0, \infty)$ is measurable, then*

$$(24) \quad \left(\int_{\mathbb{R}^n} g(x)^{n/(n-s)} dx\right)^{(n-s)/n} \leq \int_0^\infty |\{g \geq t\}|^{(n-s)/n} dt.$$

If the right side is finite, then there is equality precisely if $g = c 1_E$ for some $E \subset \mathbb{R}^n$ of finite measure and $c \geq 0$.

Proof. Let the right side of (24) be finite. By Fubini's theorem, we have

$$\int_{\mathbb{R}^n} g(x)^{n/(n-s)} dx = \frac{n}{n-s} \int_0^\infty t^{s/(n-s)} |\{g \geq t\}| dt.$$

Since $r \mapsto |\{g \geq r\}|$ is monotone decreasing, we obtain for $t > 0$,

$$\begin{aligned} t^{s/(n-s)} |\{g \geq t\}| &= (t |\{g \geq t\}|^{(n-s)/n})^{s/(n-s)} |\{g \geq t\}|^{(n-s)/n} \\ &\leq \left(\int_0^t |\{g \geq r\}|^{(n-s)/n} dr \right)^{s/(n-s)} |\{g \geq t\}|^{(n-s)/n} \\ &= \frac{n-s}{n} \frac{d}{dt} \left(\int_0^t |\{g \geq r\}|^{(n-s)/n} dr \right)^{n/(n-s)}. \end{aligned}$$

Hence

$$\left(\int_{\mathbb{R}^n} g(x)^{n/(n-s)} dx \right)^{(n-s)/n} \leq \int_0^\infty |\{g \geq t\}|^{(n-s)/n} dt$$

and there is equality precisely if $g = c 1_E$ for some $E \subset \mathbb{R}^n$ of finite measure and $c \geq 0$. \square

Proof of Theorem 1. First, assume that f is non-negative. By the co-area formula (12) and (14),

$$\begin{aligned} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{\|x - y\|_K} dx dy &= 2 \int_0^\infty P_s(\{f \geq t\}, K) dt \\ &= n \int_0^\infty \tilde{V}_{-s}(K, \Pi^{*,s}\{f \geq t\}) dt. \end{aligned}$$

Hence, by the dual mixed volume inequality (9) and (14),

$$\begin{aligned} |\Pi^{*,s} f| &= \int_0^\infty \tilde{V}_{-s}(\Pi^{*,s} f, \Pi^{*,s}\{f \geq t\}) dt \\ &\geq |\Pi^{*,s} f|^{(n+s)/n} \int_0^\infty |\Pi^{*,s}\{f \geq t\}|^{-s/n} dt \end{aligned}$$

and

$$|\Pi^{*,s} f|^{-s/n} \geq \int_0^\infty |\Pi^{*,s}\{f \geq t\}|^{-s/n} dt.$$

The equality cases of inequality (1) and of the dual mixed volume inequality, applied to (21) for $K = B^n$, show that

$$\alpha_{n,s} n \omega_n^{(n+s)/n} = |\Pi^{*,s} B^n|^{s/n} |B^n|^{(n-s)/n}.$$

By the fractional Petty projection inequality from Theorem 16, it now follows that

$$\begin{aligned} |\Pi^{*,s} f|^{-s/n} &\geq \frac{|\Pi^{*,s} B^n|^{-s/n}}{|B^n|^{(n-s)/n}} \int_0^\infty |\Pi^{*,s}\{f \geq t\}|^{-s/n} dt \\ &\geq \frac{1}{\alpha_{n,s} n \omega_n^{(n+s)/n}} \int_0^\infty |\{f \geq t\}|^{(n-s)/s} dt. \end{aligned}$$

In particular, the last term is finite. We apply Lemma 18 to obtain

$$(25) \quad \alpha_{n,s} n \omega_n^{(n+s)/n} |\Pi^{*,s} f|^{-s/n} \geq \|f\|_{\frac{n}{n-s}}.$$

Combining the equality cases of the fractional Petty projection inequality and Lemma 18, we obtain the equality case for non-negative f .

For general f and $x, y \in \mathbb{R}^n$, we use $|f(x) - f(y)| \geq ||f(x)| - |f(y)||$, where equality holds if and only if $f(x)$ and $f(y)$ are both non-negative or non-positive. Applying this inequality in the definition of $\Pi^{*,s}f$ we obtain

$$|\Pi^{*,s}|f||^{-s/n} \leq |\Pi^{*,s}f|^{-s/n}$$

with equality if and only if f has constant sign for almost every $x, y \in \mathbb{R}^n$. Using (25) for $|f|$, we obtain the first inequality of the theorem and its equality case.

For the second inequality, we set $K = B^n$ in (14) and apply the dual mixed volume inequality (9) to obtain

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|}{|x - y|^{n+s}} dx dy = n\tilde{V}_{-s}(B^n, \Pi^{*,s}f) \geq n\omega_n^{(n+s)/n} |\Pi^{*,s}f|^{-s/n}.$$

There is equality precisely if $\Pi^{*,s}f$ is a ball, which is the case for radially symmetric functions. \square

9. FRACTIONAL ZHANG PROJECTION INEQUALITIES AND RADIAL MEAN BODIES

Let $E \subset \mathbb{R}^n$ be a convex body. For $p > -1$, Gardner and Zhang [13] defined the radial p -th mean body of E , by its radial function for $\xi \in \mathbb{S}^{n-1}$, as

$$\rho_{R_p E}(\xi)^p = \frac{1}{|E|} \int_E \rho_{E-x}(\xi)^p dx$$

for $p \neq 0$ and as

$$\log(\rho_{R_0 E}(\xi)) = \frac{1}{|E|} \int_E \log(\rho_{E-x}(\xi)) dx.$$

They showed that $R_p E$ is a star body for $p > -1$ and a convex body for $p \geq 0$.

Let $0 < s < 1$. For a star body $K \subset \mathbb{R}^n$ and a convex body $E \subset \mathbb{R}^n$, we obtain from Fubini's theorem that

$$\begin{aligned} \int_E \int_{E^c} \frac{1}{\|x - y\|_K^{n+s}} dx dy &= \int_E \int_{(E-x)^c} \frac{1}{\|z\|_K^{n+s}} dz dx \\ (26) \qquad \qquad \qquad &= \frac{1}{s} \int_E \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n+s} \rho_{E-x}(\xi)^{-s} d\xi dx \\ &= \frac{|E|}{s} \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n+s} \rho_{R_{-s}E}(\xi)^{-s} d\xi. \end{aligned}$$

Hence, by (21),

$$\tilde{V}_{-s}(K, \Pi^{*,s}E) = \frac{2|E|}{s} \tilde{V}_{-s}(K, R_{-s}E)$$

for every star body K . By the equality case of the dual mixed volume inequality (9), it follows that

$$\Pi^{*,s}E = \left(\frac{s}{2|E|} \right)^{\frac{1}{s}} R_{-s}E$$

for every convex body $E \subset \mathbb{R}^n$.

Hence, we can reformulate inequalities obtained by Gardner and Zhang [13] to obtain inequalities for fractional polar projection bodies. In particular, Theorem 5.5 and Lemma 5.7 from [13] imply that

$$\left(\frac{|\Pi^{*,s}E|}{|\Pi^{*,s}\Delta|}\right)^{-s/n} \leq \left(\frac{|E|}{|\Delta|}\right)^{(n-s)/n}$$

for $E \subset \mathbb{R}^n$ a convex body and $0 < s < 1$ with equality precisely if E is a simplex, where Δ is any n -dimensional simplex in \mathbb{R}^n . Letting $s \rightarrow 1^-$ and using Theorem 3, we obtain the Zhang projection inequality [37] (without the equality case), which was reproved in [13]:

$$\left(\frac{|\Pi^*E|}{|\Pi^*\Delta|}\right)^{-1/n} \leq \left(\frac{|E|}{|\Delta|}\right)^{(n-1)/n}$$

for $E \subset \mathbb{R}^n$ a convex body.

Conversely, we can reformulate Theorem 16 for radial mean bodies and obtain sharp affine isoperimetric inequalities for radial p -th mean bodies for $-1 < p < 0$. Using also a variation of our approach to Theorem 16, we obtain the following result, which proves a natural conjecture in the field. We have included the case $p = n$ from [13, Lemma 5.7] in the statement of the result.

Theorem 19. *If $E \subset \mathbb{R}^n$ is a convex body, then*

$$(27) \quad \begin{aligned} \frac{|\mathbf{R}_p E|}{|E|} &\geq \frac{|\mathbf{R}_p B^n|}{|B^n|} && \text{for } -1 < p < 0 \text{ and } p > n, \\ \frac{|\mathbf{R}_p E|}{|E|} &\leq \frac{|\mathbf{R}_p B^n|}{|B^n|} && \text{for } 0 < p < n, \end{aligned}$$

with equality if and only if E is an ellipsoid, where for $p > -1$ and $p \neq 0, n$,

$$\frac{|\mathbf{R}_p B^n|}{|B^n|} = \left(\frac{2^{p+1}\omega_{n+p}}{(p+1)\omega_n\omega_{p+1}}\right)^{n/p}.$$

Moreover,

$$\frac{|\mathbf{R}_0 E|}{|E|} = 2^n e^{\frac{n}{2}(\psi(\frac{1}{2}) - \psi(\frac{n}{2} + 1))} \quad \text{and} \quad \frac{|\mathbf{R}_n E|}{|E|} = 1$$

for every convex body $E \subset \mathbb{R}^n$, where ψ is the digamma function.

Proof. Let $K \subset \mathbb{R}^n$ be a star body. It follows from Fubini's theorem that

$$(28) \quad \begin{aligned} \int_E \int_E \frac{1}{\|x-y\|_K^{n-p}} dx dy &= \int_E \int_{E-x} \frac{1}{\|z\|_K^{n-p}} dz dx \\ &= \frac{1}{p} \int_E \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n-p} \rho_{E-x}(\xi)^p d\xi dx \\ &= \frac{|E|}{p} \int_{\mathbb{S}^{n-1}} \rho_K(\xi)^{n-p} \rho_{\mathbf{R}_p E}(\xi)^p d\xi \\ &= \frac{n|E|}{p} \tilde{V}_p(K, \mathbf{R}_p E). \end{aligned}$$

For $0 < p < n$, the Riesz rearrangement inequality, Theorem 7, implies that

$$\begin{aligned} \int_E \int_E \frac{1}{\|x-y\|_K^{n-p}} dx dy &\leq \int_{E^*} \int_{E^*} \frac{1}{\|x-y\|_{K^*}^{n-p}} dx dy \\ &= \left(\frac{|E|}{|B^n|}\right)^{(n+p)/n} \left(\frac{|K|}{|B^n|}\right)^{(n-p)/n} \int_{B^n} \int_{B^n} \frac{1}{|x-y|^{n-p}} dx dy. \end{aligned}$$

As in the proof of Theorem 9, we obtain from Theorem 8 that there is equality precisely if E is an ellipsoid. Hence, setting $K = \mathbf{R}_p E$, we obtain from (28) that

$$\begin{aligned} \frac{|\mathbf{R}_p E|}{|\mathbf{R}_p B^n|} &= \frac{p}{n|E| |\mathbf{R}_p B^n|} \int_E \int_E \frac{1}{\|x-y\|_{\mathbf{R}_p E}^{n-p}} dx dy \\ &\leq \left(\frac{|E|}{|B^n|}\right)^{p/n} \left(\frac{|\mathbf{R}_p E|}{|\mathbf{R}_p B^n|}\right)^{(n-p)/n} \end{aligned}$$

and the result follows.

For $p > n$, we choose $c > 0$ so large that $c - \|x-y\|_K^{p-n}$ is non-negative on $E \times E$. This is possible, since E is compact. We apply the Riesz rearrangement inequality and its equality case and obtain the result.

Next, we calculate the constants for $p > 0$ and $-1 < p < 0$. For $p > 0$, we obtain by (28) that

$$\left(\frac{|\mathbf{R}_p B^n|}{|B^n|}\right)^{1/n} = \left(\frac{p}{n\omega_n^2} \int_{B^n} \int_{B^n} \frac{1}{|x-y|^{n-p}} dx dy\right)^{1/p}$$

and by [33, Theorem 7.2.7 and Theorem 8.6.6] that

$$\int_{B^n} \int_{B^n} \frac{1}{|x-y|^{n-p}} dx dy = \frac{2^{p+1} n \omega_n \omega_{n+p}}{p(p+1) \omega_{p+1}}.$$

For $-1 < p < 0$, we obtain by (26) that

$$\left(\frac{|\mathbf{R}_p B^n|}{|B^n|}\right)^{1/n} = \left(\frac{-p}{n\omega_n^2} \int_{B^n} \int_{B^{nc}} \frac{1}{|x-y|^{n-p}} dx dy\right)^{1/p}$$

and obtain the result from (11).

The case $p = n$ follows from the polar coordinate formula of volume and the definition of the radial n -th mean body. To obtain the statement for $p = 0$, we take the limit as $p \rightarrow 0$ in the second equation of (27), use that $|\mathbf{R}_p E|$ depends continuously on p and obtain

$$\frac{|\mathbf{R}_0 E|}{|E|} = \frac{|\mathbf{R}_0 B^n|}{|B^n|}$$

for every convex body $E \subset \mathbb{R}^n$. By (28),

$$\left(\frac{|\mathbf{R}_p B^n|}{|B^n|}\right)^{1/n} = \left(\frac{p}{n\omega_n^2} \int_{B^n} \int_{B^n} \frac{1}{|x-y|^{n-p}} dx dy\right)^{1/p}$$

for $p > 0$. By [33, Theorem 7.2.7 and Theorem 8.6.6],

$$\int_{B^n} \int_{B^n} \frac{1}{|x-y|^{n-p}} dx dy = \frac{2^{p+1} n \omega_n \omega_{n+p}}{p(p+1) \omega_{p+1}},$$

where $\omega_q = \pi^{q/2}/\Gamma(q/2 + 1)$ for $q > 0$. We obtain

$$\left(\frac{|R_0 B^n|}{|B^n|}\right)^{1/n} = \lim_{p \rightarrow 0^+} \left(\frac{2^{p+1}\omega_{n+p}}{(p+1)\omega_n\omega_{p+1}}\right)^{1/p} = 2e^{\frac{1}{2}(\psi(\frac{1}{2}) - \psi(\frac{n}{2} + 1))},$$

which concludes the proof. \square

Acknowledgments. J. Haddad was partially supported by CNPq, reference no. PQ 305559/2021-4 and by PIADI2020, reference no. P20_00664. M. Ludwig was supported, in part, by the Austrian Science Fund (FWF): P 34446.

REFERENCES

- [1] F. J. Almgren, Jr. and E. H. Lieb, *Symmetric decreasing rearrangement is sometimes continuous*, J. Amer. Math. Soc. **2** (1989), 683–773.
- [2] K. J. Böröczky, *Stronger versions of the Orlicz-Petty projection inequality*, J. Differential Geom. **95** (2013), 215–247.
- [3] J. Bourgain, H. Brezis, and P. Mironescu, *Another look at Sobolev spaces*, In: Optimal Control and Partial Differential Equations (J. L. Menaldi, E. Rofman and A. Sulem, eds.). A volume in honor of A. Bensoussans’s 60th birthday, Amsterdam: IOS Press; Tokyo: Ohmsha, 2001.
- [4] H. J. Brascamp, E. H. Lieb, and J. M. Luttinger, *A general rearrangement inequality for multiple integrals*, J. Funct. Anal. **17** (1974), 227–237.
- [5] A. Burchard, *Cases of equality in the Riesz rearrangement inequality*, Ann. of Math. (2) **143** (1996), 499–527.
- [6] A. Cianchi, E. Lutwak, D. Yang, and G. Zhang, *Affine Moser-Trudinger and Morrey-Sobolev inequalities*, Calc. Var. Partial Differential Equations **36** (2009), 419–436.
- [7] J. Dávila, *On an open question about functions of bounded variation*, Calc. Var. Partial Differential Equations **15** (2002), 519–527.
- [8] L. Evans and R. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.
- [9] A. Figalli, N. Fusco, F. Maggi, V. Millot, and M. Morini, *Isoperimetry and stability properties of balls with respect to nonlocal energies*, Comm. Math. Phys. **336** (2015), 441–507.
- [10] G. Folland, *Real analysis*, second ed., Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, 1999, Modern techniques and their applications, A Wiley-Interscience Publication.
- [11] R. Frank and R. Seiringer, *Non-linear ground state representations and sharp Hardy inequalities*, J. Funct. Anal. **255** (2008), 3407–3430.
- [12] R. Gardner, *Geometric Tomography*, Second ed., Encyclopedia of Mathematics and its Applications, vol. 58, Cambridge University Press, Cambridge, 2006.
- [13] R. Gardner and G. Zhang, *Affine inequalities and radial mean bodies*, Amer. J. Math. **120** (1998), 505–528.
- [14] C. Haberl and F. Schuster, *General L_p affine isoperimetric inequalities*, J. Differential Geom. **83** (2009), 1–26.
- [15] J. Haddad, C. H. Jiménez, and M. Montenegro, *Sharp affine Sobolev type inequalities via the L_p Busemann-Petty centroid inequality*, J. Funct. Anal. **271** (2016), 454–473.
- [16] N. J. Kalton, N. T. Peck, and J. W. Roberts, *An F -space sampler*, London Mathematical Society Lecture Note Series, vol. 89, Cambridge University Press, Cambridge, 1984.
- [17] A. Kreuml, *The anisotropic fractional isoperimetric problem with respect to unconditional unit balls*, Commun. Pure Appl. Anal. **20** (2021), 783–799.
- [18] D. Langharst, M. Roysdon, and A. Zvavitch, *General measure extensions of projection bodies*, Proc. London Math. Soc., in press.
- [19] Y. Lin, *The Petty projection inequality for sets of finite perimeter*, Calc. Var. Partial Differential Equations **60** (2021), Paper No. 196, 18.
- [20] M. Ludwig, *Minkowski valuations*, Trans. Amer. Math. Soc. **357** (2005), 4191–4213.

- [21] M. Ludwig, *Anisotropic fractional perimeters*, J. Differential Geom. **96** (2014), 77–93.
- [22] M. Ludwig, *Anisotropic fractional Sobolev norms*, Adv. Math. **252** (2014), 150–157.
- [23] E. Lutwak, D. Yang, and G. Zhang, *L_p affine isoperimetric inequalities*, J. Differential Geom. **56** (2000), 111–132.
- [24] E. Lutwak, D. Yang, and G. Zhang, *Sharp affine L_p Sobolev inequalities*, J. Differential Geom. **62** (2002), 17–38.
- [25] E. Lutwak, D. Yang, and G. Zhang, *Optimal Sobolev norms and the L^p Minkowski problem*, Int. Math. Res. Not. (2006), Art. ID 62987, 21.
- [26] E. Lutwak, D. Yang, and G. Zhang, *Orlicz projection bodies*, Adv. Math. **223** (2010), 220–242.
- [27] V. H. Nguyen, *New approach to the affine Pólya-Szegő principle and the stability version of the affine Sobolev inequality*, Adv. Math. **302** (2016), 1080–1110.
- [28] C. M. Petty, *Isoperimetric problems*, Proceedings of the Conference on Convexity and Combinatorial Geometry (Univ. Oklahoma, Norman, Okla., 1971), Dept. Math., Univ. Oklahoma, Norman, Okla., 1971, 26–41.
- [29] C. A. Rogers, *Two integral inequalities*, J. London Math. Soc. **31** (1956), 235–238.
- [30] C. A. Rogers, *A single integral inequality*, J. London Math. Soc. **32** (1957), 102–108.
- [31] R. Schneider, *Convex Bodies: the Brunn-Minkowski Theory*, Second expanded ed., Encyclopedia of Mathematics and its Applications, vol. 151, Cambridge University Press, Cambridge, 2014.
- [32] M. Schmuckenschläger, *Petty’s projection inequality and Santaló’s affine isoperimetric inequality*, Geom. Dedicata **57** (1995), 285–295.
- [33] R. Schneider and W. Weil, *Stochastic and Integral Geometry*, Probability and its Applications (New York), Springer-Verlag, Berlin, 2008.
- [34] A. Visintin, *Nonconvex functionals related to multiphase systems*, SIAM J. Math. Anal. **21** (1990), 1281–1304.
- [35] T. Wang, *The affine Sobolev-Zhang inequality on $BV(\mathbb{R}^n)$* , Adv. Math. **230** (2012), 2457–2473.
- [36] T. Wang, *The affine Pólya-Szegő principle: equality cases and stability*, J. Funct. Anal. **265** (2013), 1728–1748.
- [37] G. Zhang, *Restricted chord projection and affine inequalities*, Geom. Dedicata **39** (1991), 213–222.
- [38] G. Zhang, *The affine Sobolev inequality*, J. Differential Geom. **53** (1999), 183–202.

DEPARTAMENTO DE MATEMÁTICA, ICEx, UNIVERSIDADE FEDERAL DE MINAS GERAIS, 30.123-970, BELO HORIZONTE, BRAZIL

Email address: `jhaddad@mat.ufmg.br`

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNIVERSITÄT WIEN, WIEDNER HAUPTSTRASSE 8-10/1046, 1040 WIEN, AUSTRIA

Email address: `monika.ludwig@tuwien.ac.at`